

APOE ϵ 4 allele predicts faster cognitive decline in mild Alzheimer disease



S. Cosentino, PhD
N. Scarmeas, MD
E. Helzner, PhD
M.M. Glymour, PhD
J. Brandt, PhD
M. Albert, PhD
D. Blacker, MD, PhD
Y. Stern, PhD

Address correspondence and reprint requests to Dr. Yaakov Stern, Columbia University Medical Center, 630 West 168th Street, P&S Box 16, New York, NY 10032
ys11@columbia.edu

ABSTRACT

Objective: To determine whether APOE ϵ 4 predicts rate of cognitive change in incident and prevalent Alzheimer disease (AD).

Methods: Individuals were recruited from two longitudinal cohort studies—the Washington Heights and Inwood Columbia Aging Project (WHICAP; population-based) and the Predictors Study (clinic-based)—and were followed for an average of 4 years. Three samples of participants diagnosed with AD, with diverse demographic characteristics and baseline cognitive functioning, were studied: 1) 199 (48%) of the incident WHICAP cases; 2) 215 (54%) of the prevalent WHICAP cases; and 3) 156 (71%) of the individuals diagnosed with AD in the Predictors Study. Generalized estimating equations were used to test whether rate of cognitive change, measured using a composite cognitive score in WHICAP and the Mini-Mental State Examination in Predictors, varied as a function of ϵ 4 status in each sample.

Results: The presence of at least one ϵ 4 allele was associated with faster cognitive decline in the incident population-based AD group ($p = 0.01$). Parallel results were produced for the two prevalent dementia samples only when adjusting for disease severity or excluding the most impaired participants from the analyses.

Conclusion: APOE ϵ 4 may influence rate of cognitive decline most significantly in the earliest stages of Alzheimer disease. *Neurology*® 2008;70:1842-1849

GLOSSARY

AD = Alzheimer disease; **BDAE** = Boston Diagnostic Aphasia Examination; **CDR** = Clinical Dementia Rating; **DSM-III-R** = *Diagnostic and Statistical Manual of Mental Disorders*, Revised Third Edition; **ECT** = electroconvulsive treatment; **GCA** = growth curve analysis; **GEE** = generalized estimating equations; **HCFA** = Health Care Finance Administration; **MMSE** = Mini-Mental State Examination; **WHICAP** = Washington Heights and Inwood Columbia Aging Project.

The rate of cognitive deterioration during the course of Alzheimer disease (AD) varies markedly across individuals^{1,2} and is likely driven by a combination of genetic and environmental factors. Given its role in disease onset,³ the apolipoprotein E (APOE) ϵ 4 allele may contribute to differential rates of cognitive decline in AD; however, this hypothesis has been explored for over 10 years with no resolution. ϵ 4 Carriers have been found to demonstrate both slower^{4,6} and faster rates of decline,⁷⁻¹³ potentially reflecting methodologic differences in recruitment techniques, the measurement of cognition, follow-up time, and participants' stage of dementia. A subset of studies has also demonstrated non-differential rates of decline in relation to ϵ 4 status¹⁴⁻²⁴; however, most of these studies had relatively small sample sizes, which may have limited their ability to detect differential change over time.

e-Pub ahead of print on April 9, 2008, at www.neurology.org.

From the Cognitive Neuroscience Division of the Gertrude H. Sergievsky Center (S.C., N.S., Y.S.), the Taub Institute for Research in Alzheimer's Disease and the Aging Brain (N.S., Y.S., E.H.), and the Department of Neurology (N.S., Y.S.), Columbia University Medical Center, New York; Department of Epidemiology (M.M.G.), Mailman School of Public Health, Columbia University, New York, NY; the Departments of Neurology (J.B., M.A.) and Psychiatry and Behavioral Sciences (J.B., M.A.), Johns Hopkins University School of Medicine, Baltimore, MD; and the Department of Psychiatry (D.B.), Massachusetts General Hospital, Harvard Medical School, Boston.

Supported by federal grants AG0732, AG00261, RR00645, and R01AG007370.

Disclosure: Dr. Stern provides consultation to pharmaceutical companies unrelated to this study including Elan, Esai, and Wyeth. Drs. Cosentino, Scarmeas, Helzner, Glymour, Brandt, Albert, and Blacker report no conflicts of interest.

Presented in part at the 10th International Conference on AD and Related Disorders, Madrid, Spain, July 2006.

The current study was undertaken to comprehensively evaluate the effect of $\epsilon 4$ on rate of cognitive decline in three separate samples of participants with AD: incident cases from a population-based study, prevalent cases from a population-based study, and prevalent cases recruited from the clinic. Few if any prior studies have examined incident cases in which rates of cognitive decline can be observed from the earliest stages of dementia. More typically, analyses have included cases with prevalent disease frequently drawn from clinic samples, a recruitment approach that may lead to biased sampling. The simultaneous study of these three samples, which represent both incident and prevalent disease, as well as diverse demographic characteristics, recruitment techniques, and assessment procedures, allowed for one of the most comprehensive studies to date of the role of $\epsilon 4$ in cognitive decline following diagnosis of AD. Although analyses of the population-based incident sample likely have the best internal and external validity, in order to understand the discrepancies in prior studies, it is valuable to repeat the analyses across an array of samples reflecting methodologic differences in prior work.

METHODS Participants. Three samples of participants with AD (population-based incident, population-based prevalent, and clinic-based) were drawn from two different research studies as follows.

Population-based study (WHICAP). The first two samples of participants with AD were incident and prevalent samples drawn from the Washington Heights and Inwood Columbia Aging Project (WHICAP), a prospective, population-based study of aging and dementia in 4,309 Medicare recipients, 65 years and older, residing in northern Manhattan (Washington Heights, Hamilton Heights, Inwood) that has been described in detail in earlier work.^{25,26} Briefly, a stratified random sample of 50% of all persons older than 65 years was obtained from the Health Care Finance Administration (HCFA). All persons were sent a letter from HCFA explaining that they had been selected to participate in a study of aging by investigators at Columbia University. Each participant underwent an in-person interview of general health and functional ability at the time of study entry followed by a standardized assessment, including medical history, physical and neurologic examination, and a neuropsychological battery. Ethnic group was classified by participant's self-report using the format of the 1990 US Census. Participants were asked if they considered themselves white, black, or other, and then were asked if they were Hispanic. Participants were recruited at two time points (1992–

1994 and 1999–2002). They have been followed at approximately 18-month intervals with similar assessments at each interval. Evaluations were conducted in either English or Spanish, based on the primary language or preference of the participant. Recruitment, informed consent, and study procedures were approved by the Institutional Review Boards of Columbia Presbyterian Medical Center and Columbia University Health Sciences and the New York State Psychiatric Institute.

Consensus diagnoses of “dementia” or “no dementia” were based on physician-administered physical and neurologic examinations in conjunction with a standardized neuropsychological battery²⁷ according to criteria outlined in the *Diagnostic and Statistical Manual of Mental Disorders*, Revised Third Edition (DSM-III-R) and determined at a consensus conference attended by neurologists and neuropsychologists. Evidence of social or occupational function deficits was required and results from the neuropsychological battery were considered. All available ancillary information, including medical charts and imaging studies, was considered in the evaluations. Medical diagnoses were assigned when applicable. The type of dementia was subsequently determined. Diagnosis of probable or possible AD was made based on criteria of the National Institute of Neurological and Communicative Disorders and Stroke–AD and Related Disorders Association.²⁸ The study was reviewed and approved by the Columbia University institutional review board, and written informed consent was obtained from all subjects.

The first sample was comprised of 199 (48%) of the 411 incident AD cases in WHICAP. All cases were mild at diagnosis defined by a score of 1 on the Clinical Dementia Rating (CDR) scale.²⁹ A total of 56 (14%) cases were excluded due to unavailable APOE genotype. A total of 156 (38%) additional cases were ineligible due to lack of follow-up cognitive data; many of these were diagnosed relatively recently and had not yet been re-evaluated despite ongoing participation in the study. The 212 incident cases from WHICAP excluded from the current study did not differ significantly from the final incident sample in terms of sex ($p = 0.08$) or cognitive score at diagnosis ($p = 0.09$). However, excluded participants were approximately 3 years older at diagnosis (mean = 84.60; $SD = 6.42$) than the final sample (mean = 81.97; $SD = 6.50$), $F(1,409) = 16.84$, $p < 0.01$, and slightly more educated (mean = 8.02; $SD = 4.79$) than the final sample (mean = 6.30; $SD = 4.50$), $F(1,407) = 13.56$, $p < 0.01$. The final sample of 199 individuals was followed for an average of 2.9 (1.3) visits (beginning with the incident visit). Approximately 20% of participants who did not die were lost to follow-up after each of the first five visits. The proportion of individuals lost to follow-up increased once individuals participated at least five times.

The second sample was comprised of 215 (54%) of the 397 prevalent AD cases in WHICAP who met criteria for mild AD (CDR 1) at baseline. A total of 120 (30%) participants were excluded due to unavailable APOE genotyping, and an additional 55 (14%) were excluded due to lack of follow-up data. The 182 prevalent cases of mild AD from WHICAP excluded from the current study did not differ significantly from the final prevalent sample in terms of sex ($p = 0.09$) or baseline cognitive score ($p = 0.47$). However, excluded participants were approximately 4 years older at baseline (mean = 84.13; $SD = 7.82$) than the final sample (mean = 80.70; $SD = 7.10$), $F(1,395) = 21.43$, $p < 0.01$, and slightly more educated (mean = 7.14; $SD = 4.30$) than the

final sample (mean = 5.90; SD = 4.20), $F(1,395) = 8.26$, $p < 0.01$. The final sample of 215 individuals was followed for an average of 3.6 (1.6) visits. Approximately 12% of participants who did not die were lost to follow-up after each of the first six visits. The proportion of individuals lost to follow-up increased once individuals participated at least six times.

Clinic-based study (Predictors II). The final sample included individuals with AD drawn from the second cohort of the Predictors Study. Recruitment for this cohort began in 1997 using methods established for the first Predictors cohort (described in detail in earlier work³⁰). Briefly, participants in the Predictors study were recruited and studied at three US sites: Columbia University, New York, Johns Hopkins University, Baltimore, and Massachusetts General Hospital/Harvard Medical School, Boston. All participants met DSM-III-R criteria for primary degenerative dementia of the Alzheimer type and National Institute of Neurological Disorders and Stroke-AD and Related Disorders Association criteria for probable AD at enrollment. Enrollment required a modified Mini-Mental State Examination (mMMSE) score of ≥ 30 (maximum = 57) which is equivalent to a score of approximately 16 or more on the Folstein MMSE.^{31,32} Exclusion criteria were diagnosis of Parkinson disease or parkinsonism at any time prior to the onset of intellectual decline, clinical or historical evidence of stroke, history of alcohol abuse or dependence, any electroconvulsive treatment (ECT) within 2 years of recruitment or 10 or more ECT sessions at any time, and history or current clinical evidence of schizophrenia or schizoaffective disorder that started before the onset of intellectual decline.

Each consecutive patient who met the criteria of the study was included, except for those who did not consent to participate or who lived too far away and were unable to return to the study site for regular follow-up. Neurologic and mental status examinations were conducted at study entry and at 6-month intervals thereafter. The cognitive function measure used for the analysis was the MMSE. Onset of illness was estimated by the treating neurologist based on discussion with the family. The study was reviewed and approved by the participating institutions' review boards, and written informed consent was obtained from all subjects.

The current study included 156 (71%) of the 221 individuals diagnosed with AD in the second cohort of the Predictors Study. A total of 62 (28%) participants from the original sample were excluded due to unavailable genotyping and 3 (1%) subjects were excluded due to lack of follow-up cognitive data. The 65 cases excluded from the current study did not differ significantly from the final sample in terms of years of education ($p = 0.09$), sex ($p = 0.11$), or baseline cognition ($p = 0.39$). However, excluded participants were approximately 3 years older at baseline (mean = 78.57; SD = 8.33) than the final sample (mean = 75.30; SD = 7.70), $F(1,219) = 7.72$, $p < 0.01$. The final sample of 156 individuals was followed for an average of 3.6 (1.6) visits. Approximately 3% of participants who did not die were lost to follow-up after each of the first seven visits. The proportion of individuals lost to follow-up increased once individuals participated at least seven times.

Predictors. Presence of at least one $\epsilon 4$ allele was the primary predictor tested. The pattern of each subject's APOE isoforms was determined using the method of Hixson and Vernier.³³ Standard demographic variables (age at diagnosis

in the incident sample and age at first visit for the other cohorts, sex, ethnicity, and years of education) were included as covariates in statistical models. We chose to use age at the first visit in prevalent cases because these cases were selected early in the disease course (CDR = 1) and age at first visit was more clearly defined than an estimate of time of disease onset.

Outcome measures. The primary outcome in this analysis was rate of cognitive decline from baseline. Baseline is defined as the diagnostic visit for the incident cases, and the point of study entry for the other two samples. For the clinic-based sample, cognition was measured with the MMSE. For the two population-based samples, a composite cognitive z-score was developed to summarize performance on a variety of tests assessing five cognitive domains.²⁷ These included the following: 1) memory: total and delayed recall of the Selective Reminding Test, and the recognition component of the Benton Visual Retention Test; 2) abstract reasoning: Wechsler Adult Intelligence Scale-Revised Similarities subtest, and the Identities and Oddities subtest of the Dementia Rating Scale; 3) visual-spatial: five selected items from the Rosen drawing test and the matching component of the Benton Visual Retention Test; 4) language: 15-item Boston Naming Test, the eight high probability items from the Repetition subtest of the Boston Diagnostic Aphasia Examination (BDAE), and the first six items of the BDAE Comprehension subtest; and 5) executive speed: average scores for phonemic fluency assessed by the Controlled Oral Word Association Test, and category fluency (Animals, Food, Clothing).

The composite cognitive measure was derived as follows: each of the above 12 raw scores were transformed into z-scores using means and standard deviations of scores from 272 controls without dementia in WHICAP with a similar distribution of age, education, and ethnicity to the patients with AD. Z-scores for individual tests were then averaged to create a z-score for each cognitive domain. If more than half of the tests in a given domain were missing, the domain score was considered missing and was excluded from the analysis. The composite score was derived by averaging the five domain scores, with missing data treated in the manner described above.

Statistical analyses. One-way analyses of variance and χ^2 tests were used to examine between-group differences across the population-based samples, as well as within-group differences across APOE genotype in each sample. Generalized estimating equations (GEE)³⁴ were used to test whether APOE $\epsilon 4$ was associated with a differential rate of cognitive change within each sample. GEE takes into account the multiple visits per subject and the fact that the characteristics of the same individual over time are likely to be correlated. The repeated measures for each subject are treated as a cluster. GEE models in each sample examined the extent to which time (years from diagnosis in the incident sample and years from baseline in the prevalent cases), $\epsilon 4$ status (present or absent), and the time \times $\epsilon 4$ interaction predicted global cognition. A significant time effect would indicate that global cognition changed significantly over time in noncarriers (the reference group). A significant $\epsilon 4$ effect would indicate that cognitive performance at baseline (diagnosis for the incident sample and study entry for there remaining two samples) varied as a function of $\epsilon 4$ status. Finally, a significant inter-

Table 1 Demographic and clinical data by $\epsilon 4$ status in population and clinic Alzheimer disease samples

	WHICAP: Population incident			WHICAP: Population prevalent			Predictors II: Clinic-based		
	$\epsilon 4$	$-\epsilon 4$	All	$\epsilon 4$	$-\epsilon 4$	All	$\epsilon 4$	$-\epsilon 4$	All
Years followed	3.9 (2.8)	3.9 (2.4)	3.9 (2.5)*	4.4 (2.9)	5.2 (3.3)	5.0 (3.1)*	3.2 (1.7)	3.1 (1.5)	3.2 (1.6)
Visits	2.9 (1.3)	2.9 (1.2)	2.9 (1.3)*	3.4 (1.6)	3.8 (1.7)	3.6 (1.6)*	7.2 (2.8)	6.8 (2.5)	7.1 (2.6)
Baseline cognition	-1.1 (0.5)	-1.1 (0.5)	-1.0 (0.5)*	-1.3 (0.5)	-1.3 (0.6)	-1.3 (0.6)*	21.9 (3.6)	22.7 (3.5)	22.2 (3.6)
% Original sample with genotyping	NA	NA	86*	NA	NA	70*	NA	NA	71
% $\epsilon 4$	NA	NA	33	NA	NA	36	NA	NA	56
% $\epsilon 4$ homozygous	NA	NA	3	NA	NA	5	NA	NA	11
Baseline age	81.6 (6.2)	82.1 (6.7)	82.0 (6.5)	81.3 (7.1)	80.3 (7.1)	80.7 (7.1)	73.7 (7.5)*	77.4 (7.6)*	75.3 (7.7)
Education, y	6.4 (4.9)	6.3 (4.3)	6.3 (4.5)	5.9 (4.2)	5.9 (4.3)	5.9 (4.2)	15.0 (3.0)	14.3 (3.5)	14.7 (3.3)
% Women	79	71	73	79	74	76	55	61	58
% Caucasian	5	9	8	7	12	10	91	97	94
% Hispanic	59	62	61	61	62	62	0	0	0
% AA	36	28	31	33	26	28	8	1	5
% HTN	62	66	65	73	63	67	32	37	34
% Diabetes	27	23	24	18	24	21	8	8	8
% Stroke	23	13	17	18	19	19	NA	NA	NA

Mean (SD) are reported for continuous data. Years followed and number of visits were calculated from the first study visit in prevalent cases and the diagnostic visit in incident cases. Baseline = First visit for prevalent cases and diagnostic visit for incident cases. Baseline cognition is Mini-Mental State Examination in the clinic sample and composite cognitive score in WHICAP. History of vascular risk factors was assessed at baseline.

*Significant between sample differences across WHICAP prevalent and incident cases ($p < 0.05$).

†Significant within sample differences across $\epsilon 4$ group.

WHICAP = Washington Heights and Inwood Columbia Aging Project; NA = not applicable; AA = African American; HTN = hypertension.

action term would indicate that rate of cognitive decline varied as a function of $\epsilon 4$ status. To evaluate the contribution of other variables potentially related to cognitive change, GEE models also included demographic variables such as age at diagnostic visit, sex, years of education, and ethnicity as covariates. Supplementary analyses were conducted to examine the effect of baseline disease severity on $\epsilon 4$ -related decline, and to investigate potential differences in $\epsilon 4$ -related decline across demographically stratified groups.

RESULTS Descriptive statistics. Table 1 outlines follow-up, demographic, and clinical data in each of the three samples.

GEE: Main effects. Table 2 details the predictive value of time, $\epsilon 4$ status, and the time \times $\epsilon 4$ interaction in each of the three samples using several models. As expected, there was a significant time effect in all models, indicating that cognitive scores declined over time in the noncarriers. There was no main effect of the $\epsilon 4$ allele on baseline performance in any model.

GEE: Rate of change by APOE genotype. Incident sample. Demographically adjusted models reveal that cognitive scores in $\epsilon 4$ carriers declined more quickly than in noncarriers by an additional 6% of a z-score per year ($p = 0.01$) (figure).

Prevalent sample. Rate of cognitive change did not differ significantly as a function of $\epsilon 4$ status in

this sample. We thus hypothesized that the relationship between $\epsilon 4$ and rate of cognitive decline may attenuate after greater disease duration or severity of illness. When we restricted analyses to subjects with cognitive scores above the 50th percentile at baseline, there was a significant interaction between time and $\epsilon 4$ comparable to that seen in the incident sample. In contrast, when analyses were restricted to subjects with cognitive scores below the 50th percentile at baseline, $\epsilon 4$ was unrelated to rate of decline in the prevalent samples (table 2).

Clinic sample. Results for all GEE models were comparable to those reported for the prevalent population-based sample such that rate of cognitive change did not differ significantly as a function of $\epsilon 4$ status until analyses were restricted to subjects with cognitive scores above the 50th percentile at baseline. However, in this sample, we were also able to adjust the primary GEE model for duration of illness ($\epsilon 4$ carriers: mean [SD] = 4.25 [2.21] years; noncarriers: mean [SD] = 3.84 [2.12] years, $p = 0.25$), which revealed that $\epsilon 4$ carriers declined an additional 1.4 points on the MMSE than noncarriers per year ($p < 0.01$). This statistical correction was possible only in the clinic-based sample in

Table 2 Generalized estimating equation models examining the predictive value of $\epsilon 4$ for rate of global cognitive decline in three Alzheimer disease samples

	WHICAP: Population incident		WHICAP: Population prevalent		Predictors II: Clinic-based	
	β	p	β	p	β	p
Demographic adjusted						
Time	-0.05	<0.01	-0.04	<0.01	-10.4	<0.01
$\epsilon 4$	0.11	0.16	<0.01	0.96	-0.61	0.38
$\epsilon 4 \times \text{time}$	-0.06	0.01	-0.02	0.39	-0.58	0.12
High baseline cognition						
Time	-0.08	0.00	-0.05	0.00	-10.1	0.00
$\epsilon 4$	0.07	0.42	0.00	0.98	-0.05	0.94
$\epsilon 4 \times \text{time}$	-0.07	0.02	-0.06	0.04	-10.0	0.03
Low baseline cognition						
Time	-0.02	0.33	-0.03	0.04	-20.2	0.00
$\epsilon 4$	0.12	0.16	-0.02	0.85	0.28	0.71
$\epsilon 4 \times \text{time}$	-0.06	0.04	0.00	0.95	0.38	0.47

All adjusted model includes age at diagnosis (baseline age for prevalent cases), sex, and education. Models in the population samples are also adjusted for African American and Hispanic ethnicities. β values are not directly comparable across the clinic and population studies as the dependent variable differs across studies. Baseline = First visit for prevalent cases and diagnostic visit for incident cases. High baseline cognitive score = Analyses restricted to individuals performing in the top 50th percentile for cognitive score at baseline. Low baseline cognitive score = Analyses restricted to individuals performing in the bottom 50th percentile for cognitive score at baseline.


WHICAP = Washington Heights and Inwood Columbia Aging Project.

which the treating neurologist estimated the onset of illness based on discussion with the family.

Stratified models. Demographically stratified analyses were conducted in the incident sample to further explore the effect of $\epsilon 4$ on cognitive decline (table 3). Analyses were restricted to the incident sample in which there was a clear effect of $\epsilon 4$ on rate of decline in primary analyses. The $\epsilon 4$

allele was associated with faster rates of cognitive decline in women, Hispanic participants, Caucasian participants, and more highly educated participants. However, follow-up GEE analyses examining the interaction between these demographic variables, $\epsilon 4$ status and time, provided strong support only for faster decline in Caucasian $\epsilon 4$ carriers ($\beta = -0.46, p < 0.01$) and marginal support for faster decline in more highly educated $\epsilon 4$ carriers ($\beta = -0.12, p = 0.05$).

Figure Predicted cognitive decline by $\epsilon 4$ status in incident Alzheimer disease



Dotted line = noncarriers; solid line = $\epsilon 4$ carriers.

DISCUSSION The factors which determine rate of progression in AD remain incompletely understood. It has been proposed that carriers of the $\epsilon 4$ allele may demonstrate accelerated cognitive decline³ due to the allele's cumulative impact on beta amyloid and neurofibrillary tangle biochemical pathways.^{20,35,36} However, studies directly examining decline as a function of APOE genotype have produced markedly inconsistent findings.^{4-6,8-10,12,14,17-21} In the current study, we had a unique opportunity to examine the influence of APOE- $\epsilon 4$ on cognitive decline in both population- and clinic-based samples of participants with AD.

The population-based samples were primarily Hispanic and African American, and were on average older and less educated than the predominantly Caucasian participants in the clinic-based sample. There was also a lower proportion of $\epsilon 4$

Table 3 Stratified generalized estimating equation models examining the influence of $\epsilon 4$ for rate of decline in the incident sample

Stratification sample	No.	β	p
Young (<82 y)	101	-0.05	0.08
Old (≥ 82 y)	98	-0.07	0.05
Women	146	-0.07	<0.01
Men	53	-0.01	0.87
Hispanic	121	-0.07	0.01
African American	61	-0.02	0.68
Caucasian	15	-0.52	<0.01
Low education (≤ 6 y)	105	-0.04	0.14
High education (>6 y)	93	-0.11	<0.01

β values represent the rate of change over time as a function of $\epsilon 4$ status. Age and education groups were based on a median split. Models within a demographic category included the remaining demographic variables as covariates.

carriers in the population-based samples than in the clinic sample, consistent with the older age of the population samples,³⁷ and the fact that participants in the population samples were primarily Hispanic and African American, ethnic groups in which there may be a weaker association between $\epsilon 4$ and AD onset leading to an underrepresentation of the $\epsilon 4$ allele in participants with dementia.³⁸ Overall, however, rates of $\epsilon 4$ in all samples were comparable to other studies of AD and higher than rates in the general population.³⁸

The influence of $\epsilon 4$ on rate of cognitive decline varied across samples. In the incident sample, we were able to examine rate of change from the earliest stages of clinically observable AD. In this sample, the presence of the $\epsilon 4$ allele was associated with more rapid cognitive decline, even after adjustment for potential demographic contributors to disease course, suggesting a robust relationship between the $\epsilon 4$ allele and faster cognitive decline in incident AD. In contrast, $\epsilon 4$ was not associated with rate of change in either of the prevalent AD samples (population- or clinic-based). As a possible explanation for this inconsistency, we hypothesized that more advanced illness among prevalent cases may have limited our ability to detect $\epsilon 4$ -associated decline. In fact, when we accounted for disease severity by adjusting for disease duration or by restricting analyses to those with better cognitive function at baseline, we were able to demonstrate faster decline in $\epsilon 4$ carriers in both prevalent samples.

Our interpretation of the overall pattern of results is that $\epsilon 4$ influences cognitive decline most clearly in the earliest stages of disease and less so,

or not at all, in the moderate to severe stages. There may be several explanations for this. First, to the extent that our cognitive batteries are less sensitive to change as the disease progresses, we may be unable to detect a differential role for $\epsilon 4$ on rate of change in the later stages. It is possible that a different cognitive assessment tool or a functional measure would better capture differential rates of decline further into the disease. A second possibility is that the influence of the $\epsilon 4$ allele may be time limited, until some neurobiological threshold is crossed and after which medical, social, or disease-related variables other than $\epsilon 4$ become more prominent in determining rate of decline.

Investigations documenting faster rates of decline in $\epsilon 4$ carriers^{8-10,12,13} included two of the largest studies to date. Convincingly, a recent study detected faster rates of decline using both a linear and nonlinear mixed-effect statistical model.⁸ Faster rates of cognitive decline in $\epsilon 4$ carriers is consistent with the well established role of APOE $\epsilon 4$ as a risk factor for developing AD,^{35,39} and parallels findings from imaging studies suggesting accelerated rates of neurodegenerative change in $\epsilon 4$ carriers.^{40,41}

Three studies, including an early study from our group, found slower decline among $\epsilon 4$ carriers than noncarriers. The earliest study was problematic in that it retrospectively estimated baseline MMSE score based on age and education predicted values.⁴ The remaining studies used growth curve analysis (GCA) to determine rate of progression; however, the GCA study from our group likely had selective attrition as APOE genotyping was done 6 years after the study began. It is thus possible that $\epsilon 4$ carriers progressed more quickly in the early years of the study, and were more likely to have been lost to follow-up by the time $\epsilon 4$ data were collected. Additionally, as genotyping was completed on 42% of participants in the first cohort compared with 71% in the second (current) cohort, the current study better represents the population of $\epsilon 4$ carriers. In fact, re-examination of the Predictors I data with additional follow-up data using the GEE model described in the current article found no relationship between $\epsilon 4$ status and rate of decline in that cohort (data not shown).

In the current study, stratified analyses revealed that $\epsilon 4$ influenced rate of change differentially across a number of demographic subgroups. Specifically, $\epsilon 4$ appears to influence rate of decline primarily in Caucasians rather than Hispanics and African Americans. This pattern of results is largely consistent with the literature examining

the risk for developing AD³⁸ although there is some evidence for $\epsilon 4$ as a factor in Hispanic^{42,43} and African American populations as well.⁴⁴

APOE status may also interact with educational level; results offered marginal support for the $\epsilon 4$ allele predicting faster decline in those with higher education (greater than 6 years) only. The relevance of this finding is unclear, and further work is needed to more thoroughly investigate this potential interaction. One possibility is that $\epsilon 4$ exacerbates the increased rate of decline previously observed in individuals with higher education.²⁵

Although initial stratified analyses revealed a differential effect of $\epsilon 4$ by sex, follow-up GEE analyses did not support this finding. A sex-specific role of $\epsilon 4$ has been noted in previous studies reporting increased $\epsilon 4$ -related entorhinal atrophy in early AD most prominently in women,⁴⁵ and increased risk for age-related cognitive decline in women carrying the $\epsilon 4$ allele.⁴⁶ Studies examining the development of late-onset familial AD have also revealed a sex-associated role of APOE genotype with the $\epsilon 4$ allele conferring greater risk in women.⁴⁷ In contrast, a recent study linked $\epsilon 4$ to mortality rate in men only⁴⁸; however, the processes contributing to disease development and rate of decline early in the disease course may differ from those contributing to mortality at a later stage. Future work is needed to directly examine this potential discrepancy.

The current study was limited by the fact that $\epsilon 4$ genotyping was not available for 100% of individuals, introducing the possibility that participants do not accurately represent the distribution of $\epsilon 4$ in the population. Individuals excluded from the study were largely similar to the analysis sample with regard to basic demographic and clinical variables, except that excluded individuals were generally older and more highly educated than participants in each of the final samples. A third limitation to the current study may be the use of a linear rather than nonlinear approach to model cognitive decline, a technique that may have restricted our ability to observe differential rates of decline in the more impaired participants.

ACKNOWLEDGMENT

The authors thank Dr. Nicole Schupf for review of this manuscript and contributions.

Received July 5, 2007. Accepted in final form October 22, 2007.

REFERENCES

1. Hui JS, Wilson RS, Bennett DA, Bienias JL, Gilley DW, Evans DA. Rate of cognitive decline and mortality in Alzheimer's disease. *Neurology* 2003;61:1356–1361.
2. Storandt M, Grant EA, Miller JP, Morris JC. Rates of progression in mild cognitive impairment and early Alzheimer's disease. *Neurology* 2002;59:1034–1041.
3. Saunders AM, Strittmatter WJ, Schmechel D, et al. Association of apolipoprotein E allele epsilon 4 with late-onset familial and sporadic Alzheimer's disease. *Neurology* 1993;43:1467–1472.
4. Frisoni GB, Govoni S, Geroldi C, et al. Gene dose of the epsilon 4 allele of apolipoprotein E and disease progression in sporadic late-onset Alzheimer's disease. *Ann Neurol* 1995;37:596–604.
5. Stern Y, Brandt J, Albert M, et al. The absence of an apolipoprotein epsilon4 allele is associated with a more aggressive form of Alzheimer's disease. *Ann Neurol* 1997;41:615–620.
6. Hoyt BD, Massman PJ, Schatschneider C, Cooke N, Doody RS. Individual growth curve analysis of APOE epsilon 4-associated cognitive decline in Alzheimer disease. *Arch Neurol* 2005;62:454–459.
7. Marra C, Bizzarro A, Daniele A, et al. Apolipoprotein E epsilon4 allele differently affects the patterns of neuropsychological presentation in early- and late-onset Alzheimer's disease patients. *Dement Geriatr Cogn Disord* 2004;18:125–131.
8. Martins CA, Oulhaj A, de Jager CA, Williams JH. APOE alleles predict the rate of cognitive decline in Alzheimer disease: a nonlinear model. *Neurology* 2005; 65:1888–1893.
9. Dal Forno G, Rasmusson DX, Brandt J, et al. Apolipoprotein E genotype and rate of decline in probable Alzheimer's disease. *Arch Neurol* 1996;53:345–350.
10. Craft S, Teri L, Edland SD, et al. Accelerated decline in apolipoprotein E-epsilon4 homozygotes with Alzheimer's disease. *Neurology* 1998;51:149–153.
11. Wilson RS, Bennett DA, Bienias JL, et al. Cognitive activity and incident AD in a population-based sample of older persons. *Neurology* 2002;59:1910–1914.
12. Hirano N, Hashimoto M, Yasuda M, Kazui H, Mori E. Accelerated memory decline in Alzheimer's disease with apolipoprotein epsilon4 allele. *J Neuropsychiatry Clin Neurosci* 2003;15:354–358.
13. Kanai M, Shizuka M, Urakami K, et al. Apolipoprotein E4 accelerates dementia and increases cerebrospinal fluid tau levels in Alzheimer's disease. *Neurosci Lett* 1999;267:65–68.
14. Jonker C, Schmand B, Lindeboom J, Havekes LM, Launer LJ. Association between apolipoprotein E epsilon4 and the rate of cognitive decline in community-dwelling elderly individuals with and without dementia. *Arch Neurol* 1998;55:1065–1069.
15. Murphy GM Jr, Taylor J, Kraemer HC, Yesavage J, Tinklenberg JR. No association between apolipoprotein E epsilon 4 allele and rate of decline in Alzheimer's disease. *Am J Psychiatry* 1997;154:603–608.
16. Holmes C, Levy R, McLoughlin DM, Powell JF, Lovestone S. Apolipoprotein E: non-cognitive symptoms and cognitive decline in late onset Alzheimer's disease. *J Neurol Neurosurg Psychiatry* 1996;61:580–583.
17. Kurz A, Egensperger R, Haupt M, et al. Apolipoprotein E epsilon 4 allele, cognitive decline, and deteriora-

- tion of everyday performance in Alzheimer's disease. *Neurology* 1996;47:440–443.
18. Basun H, Grut M, Winblad B, Lannfelt L. Apolipoprotein epsilon 4 allele and disease progression in patients with late-onset Alzheimer's disease. *Neurosci Lett* 1995;183:32–34.
 19. Growdon JH, Locascio JJ, Corkin S, Gomez-Isla T, Hyman BT. Apolipoprotein E genotype does not influence rates of cognitive decline in Alzheimer's disease. *Neurology* 1996;47:444–448.
 20. Gomez-Isla T, West HL, Rebeck GW, et al. Clinical and pathological correlates of apolipoprotein E epsilon 4 in Alzheimer's disease. *Ann Neurol* 1996;39:62–70.
 21. Kleiman T, Zdanys K, Black B, et al. Apolipoprotein E epsilon4 allele is unrelated to cognitive or functional decline in Alzheimer's disease: retrospective and prospective analysis. *Dement Geriatr Cogn Disord* 2006;22:73–82.
 22. Asada T, Kariya T, Yamagata Z, Kinoshita T, Asaka A. ApoE epsilon 4 allele and cognitive decline in patients with Alzheimer's disease. *Neurology* 1996;47:603.
 23. Farlow MR, Cyrus PA, Nadel A, Lahiri DK, Brashear A, Gulanski B. Metrifonate treatment of AD: influence of APOE genotype. *Neurology* 1999;53:2010–2016.
 24. Slooter AJ, Houwing-Duistermaat JJ, van Harskamp F, et al. Apolipoprotein E genotype and progression of Alzheimer's disease: the Rotterdam Study. *J Neurol* 1999;246:304–308.
 25. Scarmeas N, Albert SM, Manly JJ, Stern Y. Education and rates of cognitive decline in incident Alzheimer's disease. *J Neurol Neurosurg Psychiatry* 2006;77:308–316.
 26. Stern Y, Gurland B, Tatemichi TK, Tang MX, Wilder D, Mayeux R. Influence of education and occupation on the incidence of Alzheimer's disease. *JAMA* 1994;271:1004–1010.
 27. Stern Y, Andrews H, Pittman J, et al. Diagnosis of dementia in a heterogeneous population. Development of a neuropsychological paradigm-based diagnosis of dementia and quantified correction for the effects of education. *Arch Neurol* 1992;49:453–460.
 28. McKhann G, Drachman D, Folstein M, Katzman R, Price D, Stadlan EM. Clinical diagnosis of Alzheimer's disease: report of the NINCDS-ADRDA Work Group under the auspices of Department of Health and Human Services Task Force on Alzheimer's Disease. *Neurology* 1984;34:939–944.
 29. Morris JC, Edland S, Clark C, et al. The consortium to establish a registry for Alzheimer's disease (CERAD). Part IV. Rates of cognitive change in the longitudinal assessment of probable Alzheimer's disease. *Neurology* 1993;43:2457–2465.
 30. Stern Y, Folstein M, Albert M, et al. Multicenter study of predictors of disease course in Alzheimer disease (the "predictors study"). I. Study design, cohort description, and intersite comparisons. *Alzheimer Dis Assoc Disord* 1993;7:3–21.
 31. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state": A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* 1975;12:189–198.
 32. Stern Y, Sano M, Paulson J, Mayeux R. Modified mini-mental state examination: validity and reliability. *Neurology* 1987;37(suppl 1):179.
 33. Hixson JE. Apolipoprotein E polymorphisms affect atherosclerosis in young males. Pathobiological Determinants of Atherosclerosis in Youth (PDAY) Research Group. *Arterioscler Thromb* 1991;11:1237–1244.
 34. Liang KY, Zeger SL. Longitudinal data analysis using generalized linear models. *Biometrika* 1986;73:13–22.
 35. Strittmatter WJ, Saunders AM, Schmechel D, et al. Apolipoprotein E: high-avidity binding to beta-amyloid and increased frequency of type 4 allele in late-onset familial Alzheimer disease. *Proc Natl Acad Sci USA* 1993;90:1977–1981.
 36. Ohm TG, Kirca M, Bohl J, Scharnagl H, Gross W, Marz W. Apolipoprotein E polymorphism influences not only cerebral senile plaque load but also Alzheimer-type neurofibrillary tangle formation. *Neuroscience* 1995;66:583–587.
 37. Meyer MR, Tschanz JT, Norton MC, et al. APOE genotype predicts when—not whether—one is predisposed to develop Alzheimer disease. *Nat Genet* 1998;19:321–322.
 38. Mayeux R. Apolipoprotein E, Alzheimer disease, and African Americans. *Arch Neurol* 2003;60:161–163.
 39. Corder EH, Saunders AM, Strittmatter WJ, et al. Gene dose of apolipoprotein E type 4 allele and the risk of Alzheimer's disease in late onset families. *Science* 1993;261:921–923.
 40. Mori E, Lee K, Yasuda M, et al. Accelerated hippocampal atrophy in Alzheimer's disease with apolipoprotein E epsilon4 allele. *Ann Neurol* 2002;51:209–214.
 41. Lehtovirta M, Kuikka J, Helisalmi S, et al. Longitudinal SPECT study in Alzheimer's disease: relation to apolipoprotein E polymorphism. *J Neurol Neurosurg Psychiatry* 1998;64:742–746.
 42. Olarte L, Schupf N, Lee JH, et al. Apolipoprotein E epsilon4 and age at onset of sporadic and familial Alzheimer disease in Caribbean Hispanics. *Arch Neurol* 2006;63:1586–1590.
 43. Rippon GA, Tang MX, Lee JH, Lantigua R, Medrano M, Mayeux R. Familial Alzheimer disease in Latinos: interaction between APOE, stroke, and estrogen replacement. *Neurology* 2006;66:35–40.
 44. Murrell JR, Price B, Lane KA, et al. Association of apolipoprotein E genotype and Alzheimer disease in African Americans. *Arch Neurol* 2006;63:431–434.
 45. Juottonen K, Lehtovirta M, Helisalmi S, Riekkinen P, Soinen H. Major decrease in the volume of the entorhinal cortex in patients with Alzheimer's disease carrying apolipoprotein E epsilon4 allele. *J Neurol Neurosurg Psychiatry* 1998;65:322–327.
 46. Mortensen EL, Hogh P. A gender difference in the association between APOE genotype and age-related cognitive decline. *Neurology* 2001;57:89–95.
 47. Martinez M, Champion D, Brice A, et al. Apolipoprotein E epsilon4 allele and familial aggregation of Alzheimer disease. *Arch Neurol* 1998;55:810–816.
 48. Dal Forno G, Carson KA, Brookmeyer R, Troncoso J, Kawas CH, Brandt J. APOE genotype and survival in men and women with Alzheimer's disease. *Neurology* 2002;58:1045–1050.