

**Objective Metamemory Testing Captures
Awareness of Deficit in Alzheimer's Disease**

Stephanie Cosentino¹, Janet Metcalfe², Brady Butterfield¹, & Yaakov Stern^{1, 2}

Cognitive Neuroscience Division of the Taub Institute,¹ Columbia University College of
Physicians and Surgeons, NY, NY

Department of Psychology², Columbia University, NY, NY

ABSTRACT

For reasons that remain unknown, there is marked inter-person variability in awareness of episodic memory loss in patients with Alzheimer's disease (AD). Existing research designs, primarily subjective in nature, have been at a relative disadvantage for evaluating disordered metamemory and its relation to the clinical and neuropathological heterogeneity of AD, as well as its prognosis for various disease outcomes. The current study sought to establish an objective means of evaluating metamemory in AD by modifying traditional metacognitive paradigms in which participants are asked to make predictions regarding their own memory performance. Variables derived from this measure were analyzed in relation to clinically rated awareness for memory loss. As predicted, a range of awareness levels existed across patients with mild to moderate AD ($n = 24$) and clinical ratings of awareness were significantly associated with verbal episodic memory monitoring ($r = .46, p = .03$). Further, patients who were rated as aware of their memory loss remained well calibrated over the course of the task whereas those rated as relatively unaware grew over-confident in their predictions $F(1, 33) = 4.19, p = .02$. Findings suggest that over-confidence may be related to impaired online error recognition and compromised use of metamemory strategies such as the Memory for Past Test heuristic. Importantly, clinically rated awareness did not vary as a function of demographic variables, global cognition, or verbal memory. However, participants characterized as relatively unaware were impaired on a nonverbal memory task as compared to aware participants, $F(1, 20) = 6.98, p = .02$. The current study provides preliminary support for the use of a recognition-based verbal episodic memory monitoring task as a quantitative measure of awareness for memory loss in AD, and offers insight into the manner in which metamemory breaks down. Discrepancies in nonverbal memory across the two awareness groups provide partial support for the idea that metamemory variability in AD may be related to the neuroanatomic presentation of the disease, with disordered awareness potentially reflective of a critical level of right hemisphere involvement.

INTRODUCTION

Episodic memory loss, the earliest and most striking signature of Alzheimer's disease (AD), is directly attributable to neuropathological changes in the hippocampus¹⁻⁴. Prerequisite for the diagnosis of AD, episodic memory loss is common to all patients. In contrast, for reasons that remain unknown, there is marked inter-person variability in awareness of this episodic deficit (metamemory)⁵⁻⁷. Disordered awareness of memory loss in dementia challenges patient safety^{8,9}, the efficacy of behavioral interventions¹⁰, and caregiver quality of life¹¹⁻¹³. Despite its frequency and clinical relevance, the etiology, nature, clinical correlates, and prognostic value of impaired awareness for memory loss remain poorly understood¹⁴⁻¹⁶. Although numerous studies have examined these issues, discrepant results across investigations highlight the complexity of the construct of awareness, and the limitations of using subjective methodological approaches to clarify the phenomenon of disordered awareness¹⁷.

Historically, disordered awareness in dementia (also referred to as anosognosia) has been characterized using clinician ratings^{6, 18-20}, or discrepancy scores that reflect differences between patients' report of their own functioning and caregivers' reports^{7, 21-24}. While these measures are informative, their subjectivity limits their ability to probe the nature and extent of disordered awareness, and to reliably assess this clinical phenomenon over time and across studies. Further, in the case of discrepancy scores, patients who endorse memory problems may still be characterized as having some degree of reduced awareness if their caregiver reports more severe problems. While caregiver impression is important, it cannot be considered an unbiased standard of accuracy, particularly given the psychological, economic, functional, and social adjustments involved with caring for a family member with dementia¹⁷. In the absence of objective methodology, we are impeded in our investigation of the etiology, components, clinical correlates, neurocognitive substrates, and prognostic value of disordered awareness in AD.

In all likelihood, disordered awareness for memory loss or other disease symptoms (anosognosia) is influenced by a combination of neurocognitive, psychological, social, and cultural variables.²⁵⁻³¹ According to Clare's biopsychosocial model, neurocognitive features likely determine the domains of unawareness (e.g., memory, executive abilities), whereas psychological variables (e.g., coping style and premorbid personality traits), and social factors (e.g., interaction with friends, family, and healthcare

professionals) contribute to the overall extent and presentation of unawareness in each individual. The complex manner in which these factors interact likely contributes to the variability in the degree and type of awareness for disease related deficits seen across individuals with AD.²⁹ Clarification of the components of awareness, such as memory monitoring abilities, may advance understanding of the construct as a whole and shed light on important aspects of disease heterogeneity.

Objective test frameworks such as Judgment of Learning (JOL) and Feeling of Knowing (FOK) are commonly used tasks³² to evaluate memory monitoring in healthy adults and have the potential to inform metamemory changes in the context of AD. Both types of judgments involve making predictions regarding future performance on specific test items related to general knowledge (semantic memory)^{33, 34} and newly learned (episodic) information.^{35, 36} Traditionally, JOLs are acquired for all items in a memory test, and evaluated in a cued recall format whereas FOKs are acquired for non-recallable items and tested in recognition format (or another context that does not rely on recall).³⁷ Both types of judgments offer the opportunity to evaluate several aspects of memory monitoring including: 1) resolution (relative accuracy), or the extent to which accuracy is high when predictions are high, and accuracy is low when predictions are low; and 2) calibration (absolute accuracy), the degree to which individuals are over or under-confident on average. Calibration can also be measured on a global level by having the participant make a judgment regarding the total number of items that he or she expects to achieve and comparing this global estimation to the total memory score. There is some evidence to suggest that on multi-trial learning tasks, healthy adults are overconfident on the first trial, and subsequently become underconfident, an effect that has been described as the underconfidence with practice (UWP) effect.³⁸ Recent work has suggested that this may be due to the use of the Memory for Past Test (MPT) heuristic to construct predictions after the first trial.³⁹ That is, predictions are based in part on performance at the previous trial rather than exclusively on current memory for items, a strategy that seems to result in predictions that underestimate accuracy.

Studies of older adults generally suggest that both semantic⁴⁰⁻⁴² and episodic memory monitoring⁴²⁻⁴⁷ are intact. An exception is Souchay and colleagues' finding that episodic FOK, but not JOL, is impaired in older adults secondary to executive dysfunction.^{48, 49} The link between FOK and executive function has previously been demonstrated in individuals with frontal lobe lesions^{50, 51} and Korsakoff's syndrome,⁵² and may reflect the fact that individuals construct FOKs with regard to non-recallable

items, a process that may place greater demands on executive abilities than JOLs (which are constructed for all items). The extent to which participants included in Souchay's study were differentially impaired in with regard to executive abilities, or whether executive changes that occur even in the course of "healthy aging" are sufficient to reduce the accuracy of FOK judgments, is unclear.

Similar to healthy elders, patients with AD generally demonstrate intact memory monitoring for general knowledge (semantic memory)^{53, 54} although this is not fully supported⁴². A more consistent finding seems to be that episodic memory monitoring is impaired in participants with AD using both JOLs⁵⁵⁻⁵⁹ and FOK⁶⁰. Most studies have used global level judgments which require participants to predict the total number of list items that they will recall, and have found that AD groups overestimate performance as compared to healthy controls. Moulin and colleagues have raised the concern that severely impaired recall in AD may hamper the interpretation of recall-based metacognitive assessment, focusing instead on evaluating metamemory processes that operate during encoding^{55, 61}. For instance, this group has demonstrated that despite being over-confident in global pre-study predictions, participants with AD generally revise post-study predictions downward⁵⁵. Additionally, participants are sensitive to objective differences in item difficulty, predicting that they are likely to recall the more difficult words. However, Moulin acknowledges that the sensitivity approach does not necessarily reflect participants' awareness of their *own* memory processes⁶¹.

Existing work has thus provided evidence that aspects of episodic memory monitoring are impaired in participants with AD in comparison to healthy elders. However, an important issue is that nearly all studies have examined AD as a homogenous group without taking into account the marked variations in clinically observed awareness of memory loss across individuals. This approach has limited investigation of the nature and extent of metamemory deficits in AD. For example, Ansell and Buck used a JOL format to evaluate the applicability of the Cognitive Awareness Model (CAM),^{14, 15} a neurocognitive model outlining three forms of anosognosia potentially applicable to AD⁵⁸. Briefly, according to the CAM, information regarding a memory failure is filtered through a comparator mechanism where the failure is compared with the personal database and recognized as an aberrant occurrence; this discrepancy is then processed in explicit awareness via the Metacognitive Awareness System. Disturbances in this overall system may result in one of three forms of anosognosia: *executive anosognosia* in which there is a failure at the "comparator level"

such that memory errors are never recognized as aberrant occurrences; *mnemonic anosognosia* in which memory errors are recognized as aberrant but are never incorporated into the “personal knowledge base”; and *primary anosognosia* in which the final “Metacognitive Awareness System” is disrupted resulting in complete absence of explicit awareness of memory or other cognitive errors.

Ansell and Bucks tested the applicability of mnemonic anosognosia in participants with AD expecting that memory errors may be recognized as failures, but are not integrated into the personal database and therefore do not influence participants’ judgments of their memory despite repeated testing. Indeed, the authors found that in comparison to healthy adults, the AD group overestimated their recall abilities across all three trials of a list-learning test. However, predictions in the AD group decreased over each trial, moving closer toward their accuracy scores, and explicit post-test ratings revealed that the AD group was aware that they had performed below predicted levels. After a delay, this awareness diminished to some extent but did not return to pre-test levels, offering only partial support for the model of mnemonic anosognosia.

A critical issue is the extent to which patient metamemory patterns differed as a function of the individual’s overall awareness of memory loss. Given the heterogeneity of awareness in AD, it is possible that relatively aware participants may have achieved metamemory scores in line with healthy elders whereas relatively unaware participants may have been less able to update their personal knowledge base over the course of the study, particularly after a delay. In fact, a study by Reed and colleagues suggested that clinical ratings of awareness based on a structured interview were moderately correlated with discrepancies between participants’ predictions and performance on a memory test⁶².

The aim of the current study was to examine metamemory heterogeneity within AD in relation to clinically rated awareness of memory loss. Comprehensive examination of memory monitoring in early AD might generate an objective assessment of awareness for memory loss that has the potential to advance our conceptualization and study of disordered awareness. Metacognitive studies in patients with AD have established the feasibility of such techniques in this population while raising important methodological issues to be considered in future studies. We evaluated memory monitoring in the context of an objective battery designed specifically for individuals with mild to moderate dementia to determine whether or not test scores would capture variations in clinically rated awareness for memory loss. We hypothesized that a range of clinically rated

levels of awareness would exist among participants, and we expected clinical ratings of awareness for memory loss to be selectively associated with memory monitoring deficits related to new learning (episodic memory) rather than general knowledge. Furthermore, clinical ratings of awareness were not expected to vary as a function of global cognition, verbal memory, or demographic variables such as age, education, ethnicity, or sex. Our overall goal was to establish a task that captured variability in awareness for episodic memory loss that might be used in future investigation of the etiology, clinical correlates and prognostic value of disordered awareness in AD.

METHODS

Participants

24 participants with AD were recruited through the Columbia University Medical Center Alzheimer's Disease Research Center and received comprehensive neurologic and neuropsychological evaluations that were reviewed in a diagnostic consensus conference attended by neurologists and neuropsychologists. Diagnoses of Alzheimer's disease were made according to the Neurologic Disorders and Stroke - Alzheimer's Disease and Related Disorders Association (NINDS-ADRDA) criteria. Given the cognitive demands of the experimental tasks and our interest in studying awareness in early AD, only patients with mild to moderate AD, defined as a score of 20 or greater on the MMSE were selected for participation. Individuals with ongoing moderate to severe psychiatric conditions were excluded from the study, as were individuals with history of head injury, stroke, and other neurologic illnesses that might impact cognition and/or the presentation of AD. All participants provided informed consent and were reimbursed \$30.00 for participation.

20 healthy elderly volunteers were recruited from senior centers in the New York City area. Participants underwent medical and neuropsychological evaluations to ensure the absence of dementia, Mild Cognitive Impairment (MCI), psychiatric illness, neurologic illness, or other conditions with cognitive sequelae.

Procedures

The study visit included examiner ratings of awareness using a modified version of the Anosognosia Rating Scale, participants' self-ratings, the MMSE, and our experimental metamemory battery. Scores for the Selective Reminding Test were obtained from the participants' most proximate neuropsychological evaluation in the

Alzheimer's Disease Research Center when visits occurred within three months of the current evaluation.

Measures

Clinical Ratings of Awareness: Testing sessions began with a brief interview to allow the examiner to make a clinical judgment regarding participant awareness of memory loss. This was done only with participants in the AD group for the purposes of establishing a clinical rating of awareness. Scores could not be assigned in healthy elders as clinical diagnosis of AD was the criteria against which self-report was measured. Examiners asked participants to discuss their opinions of their memory abilities at the current time, and assigned a score ranging from 1 to 5 on a modified version of the Anosognosia Rating Scale⁶. Participants were scored according to the following five point ordinal rating system: 5 = *Full Awareness* (Spontaneous complaint or ready admission of memory loss along with the recognition that the loss is consequential. Loss related to dementia); 4 = *Moderate Awareness* (Spontaneous admission of memory loss; however, loss is discussed in the context of "normal" age related changes. No discussion of diagnosis); 3 = *Shallow Awareness* (Inconsistent or transient recognition of memory loss, or uncertainty regarding memory loss. Patients may acknowledge inconsequential memory loss); 2 = *No Awareness* (Matter-of-fact denial of impairment in response to direct questions regarding memory loss); 1 = *Explicit Denial* (Questions regarding memory elicit vigorous denial of impairment or angry assertions of normal function). If spontaneous responses did not clearly fit into a specific rating category (e.g., "My memory is bad"), the examiner queried as appropriate (e.g., "Do you have a sense of why your memory is bad?") to extract sufficient information for assigning a score of 1-5. Responses were recorded verbatim and scored independently by two neuropsychologists (the examiner and a second rater) to determine an inter-rater reliability coefficient. The final assigned awareness score was the mean rating from the two raters.

Self-Ratings: Following examiner ratings, self-ratings were obtained using a brief anosognosia scale⁶³ that prompts participants to judge themselves on eight abilities in comparison to others their age (walk; accurately and quickly use hands and fingers; speak clearly; remember; concentrate and attend; sit still and quietly; say the word you are thinking of; and control your emotions). Participants selected one of five ratings from

a visual chart with the ratings arranged vertically (excellent; above average; average; below average; very impaired).

Mini Mental State Examination⁶⁴: This commonly used 30-item test assessing orientation, attention, language, visuospatial functioning and memory was included in the current study as a measure of global cognition.

Metamemory Battery

Training Procedure: Throughout the metamemory tasks, participants were required to make judgments about their own performance using a three point scale (Yes, Maybe, or No). In order to establish that participants could use this scale appropriately, and appreciate the difference between these judgments, we implemented a basic training procedure for this rating scale. We gave particular attention to whether or not participants appreciated the concept that “Maybe” is the best response when they cannot be sure of the answer (Will it rain three weeks from today?). The first six participants completed a brief training procedure including 6 questions about themselves (Are you seated?), the examiner (Am I eating?), and general information (Is this table red?) that required specific responses of Yes, No, or Maybe. Beginning with the seventh participant, a more comprehensive training procedure with 26 questions was implemented after a relatively impaired participant (not included in the current data set) demonstrated significant difficulty using “Maybe” appropriately. Following this training, all participants completed the three tasks described below.

1. **General Knowledge (GK):** Inclusion of the GK paradigm served two purposes. First, it acquainted participants with this test format and established that participants understood the testing framework. Second, comparable performance on this measure across awareness ratings would strengthen dissociations predicted on the Episodic JOL tasks below. The GK task consisted of thirteen general knowledge questions with a broad range of difficulty (.01 to 1.0) defined as the percentage of healthy controls able to provide the correct answer to the specific question as reported in the Nelson and Narens trivia set³⁴. The rating scale was printed on 8.5” X 11” paper, with the rating items (Yes – Maybe – No) written vertically in the center of the page, and was placed directly in front of the participant for the duration of the metamemory battery. Participants were provided with two practice items intended to acquaint them with the task, and to provide the examiner with the opportunity to correct misunderstanding of the

directions. Questions were printed on 8.5" X 11" paper with a reminder not to answer aloud at the top of each page. For example, the first practice item read, "What instrument did Miles Davis play?" After participants read the question aloud, the examiner provided the following prompt: "There are eight possible answers on the next page. Will you know which one is right – Yes, Maybe, or No?" The examiner then directed the participant to make a prediction using the rating scale. Once ratings were recorded, participants were provided with eight answer choices and asked to select the correct answer. The examiner repeated the question as necessary. Distractors were semantically related to the correct answer (e.g., trumpet, clarinet, oboe, piano, etc.). While participants were encouraged to make their best guess, "I don't know" was also accepted and scored 0 for accuracy. Once participants selected an answer choice, the examiner provided feedback regarding accuracy.

2. *Verbal Episodic Judgment of Learning (VJOL)*: In contrast to the GK metamemory test, this episodic task required participants to retain newly learned information, and to make ratings about the likelihood of recognizing each newly learned item. A recognition-based format was adopted to minimize the number of participants who performed at floor level as might occur on a recall measure. Stimuli for this task were selected to parallel stimuli in the GK task as closely as possible. Participants were asked to remember 5 pieces of factual information drawn from a subset of the Nelson and Narens³⁴ items that less than 1% of healthy adults answered correctly (e.g., "Charles Ford supposedly killed Jesse James". The examiner read the following instructions, "In the next part of the task, I will teach you eight facts. Your task is to remember this information as best you can. Then I will test your memory, giving you answer choices like before. Here are the facts". Immediately after stimuli were presented, participants were asked to predict how many of the five facts they would get right when tested in recognition format. (i.e., "Now I am going to test your memory for those facts. How many of the five do you think you will get right if I give you answer choices like before?"). These trial-based judgments enabled examination of global metamemory judgments and were used as a measurement of global calibration (over/under confidence).

Following the global judgments, participants were provided with the rating scale (Yes-Maybe-No) and the first test question. Questions were presented in the same format as in the GK task such that a reminder not to answer aloud was written at the top of the page, with the test question printed in the center of the page. The examiner

stated: “Here is the question. There are eight answer choices on the next page. Will you know which one is right?” Participants used the rating scale to make their prediction, and then chose an answer on the following page. In contrast to the GK task, feedback regarding accuracy was *not* provided throughout the VJOL task so as to enable evaluation of participants’ spontaneous adjustments to metamemory judgments over the course of the list-learning task. To control for basic familiarity effects and to ensure that participants encoded the information in a paired-associates format, the 8 answer choices included the correct response to the item, as well as 4 distractor items that are the correct answers for the remaining 4 stimuli, and 3 new distractors. The procedure was repeated three times using the same memory items. Stimuli were presented in the same order during the study phase in each trial; however, the questions and answer choices were presented in a pseudorandom order for each test trial. A delayed trial was administered after approximately 15 – 20 minutes.

3. *Nonverbal Episodic Judgment of Learning (NJOL)*: This task was designed to resemble the format of the VJOL task such that participants studied five new pieces of information over the course of four learning trials. The stimuli for this task were taken from a children’s visual memory test⁶⁵ and consisted of geometric designs located on a 4 X 2 grid of squares (5 stimuli and 3 distractors), each of which was concealed by a lid at the start of the test. The five stimuli were exposed one at a time, for five seconds each. Participants were instructed to study the design for the entire time so that they would remember the location of the design in the grid. Following study, participants were asked to estimate the number of designs out of five that they would get right (“Now I am going to test your memory for the location of the five designs. How many do you think you will get right?”). The examiner then showed the participant the first design from the stimuli book in conjunction with the rating scale, stating, “Do you think you know where this design is – Yes, Maybe, or No?” After assessing JOL and accuracy for each of the five designs, the examiner repeated this entire procedure using the same designs for trials two, three, and four. Similar to the VJOL procedure, stimuli were presented in the same order during the study phase in each trial while the prompts were presented in a pseudorandom order for each test trial. Although a delayed trial was administered after approximately 15 – 20 minutes, data are not presented in the current study since this was introduced after the start of the study and is available for only a subset of the participants.

Metamemory Scoring

Resolution: Resolution was calculated using the gamma statistic, a rank-order correlation representing the degree to which higher ratings were paired with higher accuracy scores (scored as 0 or 1) and lower ratings were paired with lower accuracy scores. In this calculation, each rating - accuracy pair is compared with the others to determine the number of concordant pairs (instances in which both the rating and accuracy are higher in one pair than another) and discordant pairs (instances in which the rating in one pair is higher than that in the second pair, but accuracy is lower). Gamma represents the number of concordant minus discordant pairs divided by the total number of discordances and concordances $(C-D)/(C+D)$. Gamma scores were calculated for each of the three metamemory tasks using all items in each task.

As the purpose of gamma is to describe the manner in which accuracy rises and falls as a function of the participant's prediction, calculation of gamma discards ties, or instances in which either rating or accuracy in one pair is equal to that in another pair. For example, all instances in which "Yes" is selected as the rating are not compared to one another in the calculation of gamma, regardless of the accuracy of the items. While this characteristic of the gamma calculation results in a "noisy" estimation of resolution, it is important for data in which many "ties" arise due to restricted range in either rating or accuracy⁶⁶. Although implementation of a scale with broader range would have offered a more precise measure of resolution, we prioritized a restricted range to ensure valid use in patients with dementia.

Item by Item Calibration: In order to calculate item-by-item calibration scores, ordinal ratings (Yes-Maybe-No) were translated into interval data (1, 0.5, and 0). Average accuracy was then subtracted from the average rating to determine the extent to which individuals were over or under confident on an item-by-item basis. A score of zero indicates perfect calibration, positive scores indicate overconfidence, and negative scores indicate underconfidence.

Global Calibration: We also calculated global calibration at each learning trial and the delay trial to evaluate individuals' ability to make pre-test predictions regarding the overall number of items that they would correctly remember out of the five studied. Global calibration scores were determined by subtracting the actual accuracy at each

trial from the predicted global accuracy, and dividing by the total number of items.

RESULTS

Demographic and cognitive variables are presented for the healthy elders and the entire AD group in Table 1, and for the two AD awareness groups in Table 2. Bivariate correlations were used within the AD group to evaluate the relationship between clinical ratings of awareness and metacognitive variables. One-way analyses of variance (ANOVAs) were used to evaluate between group differences on continuous variables. Chi-square tests were conducted to analyze the distribution of sex and ethnicity.

Awareness Ratings

An intraclass coefficient of .90 ($p < .001$) demonstrated a high degree of inter-rater reliability between the two raters applying the modified Anosognosia Rating Scale. Using the average rater score, there was a range of clinically rated awareness levels such that 8 participants were classified as Fully Aware, 7 as Moderately Aware, 6 with Shallow Awareness, 2 with No Awareness, and 1 participant who vigorously denied memory loss. Due to the relatively small sample, we created two awareness groups for certain analyses by combining the full and moderate groups into an Aware AD (AAD) group ($n=15$), and the remaining three awareness categories into an Unaware AD (UAD) group ($n=9$).

On a self-report inventory that queried participants about specific cognitive and functional abilities in comparison to others their age (1 = Very Impaired; 2 = Below Average; 3 = Average; 4 = Above Average; 5 = Excellent), ANOVA revealed that both AD groups and healthy elders rated themselves similarly for *walking, accurately and quickly using their hands and fingers, sitting still, saying the word you are thinking of, and controlling emotions*. In contrast, there were significant overall group differences ratings of the ability to *remember*, $F(2, 41) = 8.54$, $p < .01$, *speak clearly*, $F(2, 41) = 3.66$, $p = .04$, and *concentrate*, $F(2, 41) = 4.81$, $p = .01$. Planned comparisons demonstrated that the UAD group generally rated *remembering* as average ($M = 3.22$; $SD = 0.83$) and comparable to ratings by healthy elders ($M = 3.60$; $SD = 0.88$); in contrast, the AAD group assigned themselves ratings in the below average range ($M = 2.40$; $SD = .83$), significantly lower than the both the healthy elders ($p < .01$) and UAD group ($p = .03$). Self-Ratings for *speak clearly* paralleled the ratings of *remembering*,

with the AAD group assigning ratings that were significantly lower than the HE ($p = .02$) and UAD ($p = .05$) ratings. The AAD group's ratings of *concentration* were in the below average range, significantly lower than the HE ratings in the average range ($p = .00$); UAD group ratings did not differ significantly from the other groups.

Rating Scale Training Procedure

The first six participants in the study completed a six-item training procedure, and all but one participant achieved perfect scores with the final participant answering one item incorrectly. The comprehensive training for the rating scale was implemented for the remaining 18 participants (UAD = 7; AAD = 11). There was no significant difference in each group's ability to answer No, Yes, or Maybe when appropriate, $F(1, 16) = 1.88$, $p = .19$. Of note, both groups used the Maybe response correctly during training, $F(1, 16) = 1.42$, $p = .25$. Out of 13 items that required Maybe as a response, the means and standard deviations for the AAD group were 11.82 (1.60) and 12.57 (0.53) for the UAD group. These results provide evidence that participants understood the rating choices and used the scale correctly in a context that was independent of the metamemory tasks.

Experimental Task Accuracy: General Knowledge and Episodic Memory

Prior to examining metacognitive scores, ANOVAs were used to evaluate between-group difference in accuracy on three dependent variables (general knowledge, verbal episodic memory - immediate and delayed, and nonverbal episodic memory - immediate). As expected, there were significant differences on all accuracy indices between healthy elders and the AD group (Table 1). Evaluation of within-AD group differences revealed similar accuracy scores for general knowledge and verbal episodic memory; however, the UAD group scored significantly lower than the AAD group on the nonverbal episodic memory task (Table 2).

Metacognitive Scores

We conducted bivariate correlations to examine the relationship between clinical ratings of awareness and metacognitive scores (resolution and calibration) on the general knowledge, verbal episodic, and nonverbal episodic metamemory measures. In a subsequent step to establish divergent validity, metacognitive variables that correlated significantly with clinical ratings of awareness were then examined in relation to cognitive

and demographic variables.

Resolution Scores: As predicted, gamma scores on the general knowledge task were not related to clinical ratings of awareness (CRA), however, verbal episodic gamma scores were significantly and selectively correlated with CRA ($r = .46, p = .03$) such that participants who achieved higher gamma scores had higher awareness ratings (Table 3). To evaluate the extent to which disease severity might impact this association, we used a partial correlation procedure to control for MMSE. The correlation between verbal episodic gamma and clinical ratings of awareness was increased ($r = .48, p = .02$). Contrary to expectations, awareness ratings were not associated with the nonverbal episodic gamma score ($r = -.03, p = .90$). Mean gamma scores for each group were also calculated and ANOVAs with planned comparisons were used to evaluate between group differences for all metacognitive scores across the healthy elders, AAD, and UAD groups (See Table 4). In support of the above correlational results, the UAD group achieved significantly lower verbal episodic gamma scores than the AAD group.

Calibration: As predicted, calibration scores for the test of general knowledge were unrelated to clinical ratings of awareness. Contrary to predictions, neither item-level ($r = -.18, p = .44$) nor global level calibration scores ($r = -.30, p = .19$) on the verbal episodic test were related to CRA. However, follow-up analyses of between group differences revealed that, as a whole, the UAD demonstrated overconfidence on measures of global and item-level calibration scores for the verbal episodic test (See Table 4). With regard to the nonverbal episodic task, awareness ratings were significantly correlated with global calibration ($r = -.50, p = .03$) and this association persisted when the effects of disease severity (MMSE) were removed ($r = -.53, p = .02$).

Trial-Based Analyses: We conducted further analyses to examine potential changes in calibration over the course of the episodic tests. Unlike the calculation of gamma scores (which required combination of all 20 items for each episodic task), item level and global calibration scores were generated for each of the four trials in the verbal and nonverbal episodic JOL tasks. To get a general sense of calibration at the beginning and end of the episodic tests, we averaged calibration scores for the first two trials and the second two trials to represent calibration scores over the course of the test.

Repeated Measures ANOVAs with a 3 (healthy elder, aware, unaware) X 2 (Trials

1/2, Trials 3/4) design were used to evaluate the presence of interaction effects between group and trial. A significant interaction was found for item-level calibration on the verbal episodic test such that the AAD group remained evenly calibrated over time whereas the UAD group moved toward overconfidence and the healthy elders moved toward underconfidence, $F(1, 33) = 4.19, p = .02$. Planned comparisons revealed that average item-level calibration for Trials 3/4 was significantly higher in the UAD group than in the healthy elder ($p < .01$) and AAD groups ($p = .04$). There was no significant difference between scores in the latter two groups (Figure 1). Interactions were not detected for global level calibration on the verbal test, $F(1, 26) = 1.91, p = .17$, nonverbal global calibration, $F(1, 29) = 0.13, p = .88$, or nonverbal item level calibration, $F(1, 30) = 0.28, p = .76$.

Factors Underlying Metacognitive Deficits

We ran preliminary analyses to examine potential factors that may underlie deficits in resolution and calibration. One possibility is that failures in the online recognition of errors may contribute to both impaired resolution and calibration in the UAD group. We examined this by comparing the nature of errors during the verbal episodic task, specifically, the frequency with which participants chose a distractor or responded “Don’t Know” (this data was only available for the Verbal Episodic task). When controlling for total number of errors, the UAD group answered “Don’t Know” less frequently ($M = 2.33, SD = 3.01$) than the AAD group ($M = 5.33; SD = 5.61$), $F(1, 11) = 7.20, p = .01$. This data was not available for the nonverbal episodic task.

Another possible factor contributing to discrepant metamemory scores in the UAD and AAD group may have been the degree to which each awareness group implemented a metacognitive heuristic known as the Memory for Past Test (MPT) heuristic³⁹. The MPT score that we used reflects the extent to which global predictions for a given trial are related to overall accuracy on the former trial. We used one-sample *t*-tests to evaluate whether or not each group’s MPT scores ($\text{Prediction}^{\text{Trial } N} - \text{Accuracy}^{\text{Trial } N-1}$) differed significantly from zero at trials two, three, and four. Scores in the healthy elders and AAD group were not significantly different from zero at any time, suggesting that accuracy on the former trial may have guided predictions on the subsequent trial in these groups. In contrast, MPT scores in the UAD group rose significantly above zero in the final trial of the verbal episodic test and were consistently above zero throughout all trials of the nonverbal episodic test.

Although the global MPT score provides an indication of the importance of memory for past test at a cumulative level, use of the MPT heuristic is also believed to reflect the extent to which individuals make predictions for specific items on upcoming trials based on accuracy for those individual items at the previous trials. To determine whether or not the awareness groups differed in the use of this item-level approach, we calculated a backward gamma score relating item based predictions at trial N to accuracy for those specific items at trial (N-1). In fact, bivariate correlations revealed a highly significant relationship to clinical ratings of awareness ($r = .71, p < .01$) and verbal episodic gamma ($r = .74, p < .01$). Further examination of mean backward gamma scores revealed a significant difference between the aware ($M \text{ gamma} = .73, SD = .34$) and unaware groups ($M \text{ gamma} = .03, SD = .85$), $F(1, 17) = 6.89, p = .02$. This data was not available for the nonverbal episodic task.

DISCUSSION

Countless works have been dedicated to establishing a coherent understanding of syndromes of anosognosia, their etiology, and neurologic and cognitive correlates⁶⁷⁻⁷⁹. Impaired awareness for memory loss in AD represents a form of anosognosia, and may very well reflect the relative distribution of neuropathology across individuals. Disordered awareness for memory loss in AD is a clinical phenomenon that has yet to be measured in an objective framework. Historically, measurement of awareness in clinical populations has been largely subjective, and very frequently restricted to qualitative comments regarding a patient's degree of insight into the cognitive or physical sequelae related to injury or illness. The main aim of this study was to modify an objective metacognitive methodology to capture variability in clinically rated awareness of episodic memory loss in patients with AD.

Awareness Ratings

In the current study, the clinical ratings of awareness were made on the basis of a brief interview that preceded all formal testing. These ratings were compared against participants' self-ratings on a separate measure that queried regarding memory abilities in comparison to others in their peer group. Interestingly, the UAD group rated themselves as average, and comparable to healthy elders for abilities including remembering and speaking clearly, for which the AAD group provided ratings that were

significantly lower than both groups. In contrast, all three groups rated themselves as average for walking, accurately and quickly using hands and fingers, saying the word that you are thinking of, sitting still, and controlling emotions, offering some degree of specificity for the difference in memory ratings across the groups.

Metacognitive Training Procedure

We implemented a three point rating scale for metacognitive ratings. While past studies have implemented more finely graded rating scales that offer the opportunity to assess a wider range of confidence judgments,^{42, 55, 57} we felt that a three point scale would introduce fewer demands on cognitive estimation skills. This reduces the likelihood that metacognitive ratings are confounded by poor conceptualization and improper use of the rating scale. An important component of this study was to demonstrate that participants with AD are able to use the metacognitive rating scale in a valid manner, and that this did not vary as a function of awareness level. As such, we implemented a training procedure that required appropriate use of the answers Yes, No, and Maybe in response to basic questions about themselves, the examiner, and the world. All participants demonstrated the ability to use these answer choices accurately in a context independent of the metacognitive testing. This enhances our ability to interpret differential performance as a function of memory monitoring rather than non-specific cognitive factors.

Awareness Ratings and Metacognitive Scores

We evaluated clinically rated awareness in relation to two types of memory monitoring abilities, resolution and calibration. Resolution was measured with the gamma statistic, and represents the relative frequency with which higher predictions were paired with higher accuracy and lower predictions were paired with lower accuracy throughout the test. We also assessed calibration scores to evaluate over or under confidence for particular memory items as well as in the context of a global self-evaluation at each study trial.

Resolution

A main aim of this study was to demonstrate a relationship between clinically rated awareness and objective measures of episodic memory monitoring in contrast to general knowledge monitoring, global cognition, demographics, and severity of episodic

memory loss. As predicted, participants with mild to moderate AD demonstrated a range of clinically rated awareness for memory loss, and these ratings correlated significantly with gamma scores on the verbal episodic task. That is, the higher a patient's level of awareness, the more likely he or she was to adjust predictions in accord with accuracy scores. There was no relationship between general knowledge gamma scores and awareness ratings. This provides an important aspect of divergent validity, demonstrating that the association was not driven by the format of metamemory testing, but was related specifically to judgments of new learning. Further divergent validity derives from the fact that neither clinical ratings of awareness nor verbal episodic gamma scores were correlated with demographic variables, global cognition, general knowledge, or verbal memory.

To our knowledge, this is the first study to document selective differences in episodic memory monitoring as a function of clinically rated awareness for memory loss. Although Reed and colleagues demonstrated this relationship over a decade ago⁶², awareness ratings in that study were also correlated with global cognition and memory deficit. Such a pattern is not entirely surprising particularly if participants were further into the disease course than participants in the current study. As awareness for memory may worsen within individuals over time, low memory scores would likely be associated with lower awareness when studying individuals in a more advanced disease stage.

Contrary to our predictions, awareness ratings were not correlated with gamma scores for the nonverbal episodic memory task. This may be related to the fact that accuracy on the nonverbal task was significantly different across the awareness groups (a finding that is discussed below), and that gamma scores were incalculable for six participants due to a lack of variability in predictions, accuracy, or both across the 20 items. Gamma is calculated by comparing the rating and accuracy for one item with the rating and accuracy for another item. If either the rating or accuracy is "tied" with that in the second pair, the comparison does not enter into the calculation. For example, if an individual chose "Yes" as the rating for every item, gamma would be incalculable because it would be impossible to determine whether accuracy scores varied as a function of ratings. This is a limitation to the use of the gamma statistic, but can be reconciled by adding more items to the test to increase the likelihood that either ratings or accuracy will vary. We are currently revising our tasks to improve the calculability of gamma.

Calibration

Of the four calibration measures, only the global score on the nonverbal episodic task was significantly correlated with clinical ratings of awareness. However, follow up analyses revealed that this calibration score was associated with accuracy on the nonverbal memory task as well as several other measures, obscuring the extent to which this score reflects specific metamemory processes, and rendering it less compelling than the verbal episodic gamma score discussed above. In contrast to correlational analyses, between-group analyses revealed a significant difference in verbal calibration in the aware versus unaware groups with the latter group being overconfident in their predictions. Further, different patterns emerged in each group over the course of the verbal test such that the unaware group increasingly overestimated their performance whereas the aware group remained evenly calibrated and not significantly different from the healthy elders who demonstrated reduced confidence over time. This underconfidence with practice effect has been demonstrated in several previous studies and suggests that the healthy elders, and potentially the UAD group, based current predictions on accuracy from the former trial^{38, 39}. We discuss this possibility below.

Factors Underlying Metacognitive Deficits

Miscalibration is not necessarily related to resolution deficits. The fact that the UAD group demonstrates impairment in both aspects of episodic memory monitoring may implicate primary impairments in each, or a single impairment that affects both. The most parsimonious interpretation of the data is that both resolution and calibration scores are impaired secondary to a common deficit. One possibility is that the UAD group has deficient online error recognition, comparable to the executive anosognosia subtype described by the CAM. At the beginning of the verbal test, the aware and unaware groups were equally calibrated, suggesting that both groups start with a relatively accurate sense of how they will perform with only one presentation of the stimuli. However, the groups diverged in their self-assessments with repeated exposure to the stimuli. Similar to patterns seen in healthy adults³⁹, the aware group appeared to implement the Memory for Past Test heuristic (predicting that they would achieve the same amount correct as the last trial). Backward gamma scores on the verbal task suggested that they did this by assessing the accuracy of specific items from the previous trial and using this assessment to make predictions for upcoming accuracy at both the item and global level. In contrast, the unaware group did not implement these

strategies, and became over-confident over the course of the episodic test. Taken together, these results suggest that the unaware group may not recognize incoming memory failures and thus cannot accurately predict future performance; perhaps they rely on inaccurate assessments of performance, or pre-existing ideas of their memory abilities. Interestingly, after a delay, both groups are equally and accurately calibrated (similar to the first trial of the learning test). That is, when there is no trial immediately preceding the prediction (such as at trial one and at delay), the groups are indistinguishable with regard to calibration. It thus appears that calibration discrepancies in the current study occurred in direct relation to the repeated presentation of stimuli, and the extent to which each group based upcoming performance on past performance. Either failure to use the MPT strategy altogether, or use of MPT with impaired recognition of errors in the preceding trial, had a direct impact on calibration scores.

Further evidence for poor online error recognition derives from examination of errors across the verbal learning task. Specifically, when controlling for total accuracy, the UAD group made far fewer “don’t know” responses than the AAD group. That is, when the UAD group made errors, they were more likely to endorse distractors than to say “I don’t know”, a pattern of performance that may indicate that the UAD group did not recognize when they were wrong. This would lead to overconfidence in general (calibration), as well as difficulty adjusting item-by-item ratings in accord with accuracy (resolution). Although results offer some support for deficits in online error recognition, data from the current study do not clearly evaluate the degree to which participants explicitly recognize each memory error. Another possibility is that the UAD group recognized errors but were unable to incorporate these errors over time, a pattern that would be consistent with the mnemonic model of anosognosia outlined by the CAM. We are exploring these two possibilities in a follow up study in which participants are asked directly to comment on their accuracy after they provide their response. It is important to note that even within the unaware participants, there may be heterogeneity in the etiology and profile of metamemory deficits¹⁴. Overall, however, the pattern of metamemory errors seen on objective testing in the current study offers a plausible perspective regarding the syndrome of disordered awareness for memory loss seen in the daily lives of these patients.

The Neuroanatomy of Metacognitive Deficits

A broader question underlying the current line of research is the nature of the

neuropathologic substrate of self-awareness (in this case metamemory) in AD. Early Alzheimer's disease (AD) is marked by a core amnesic syndrome and heterogeneous deficits in visuospatial functioning, language abilities, and executive skills. Certainly, this disease is a progressively global illness; however, studies of the varied neuropsychological and clinical presentations of this illness have demonstrated that, within the general profile of AD, the illness is heterogeneous in its manner of presentation⁸⁰⁻⁸⁴. The various cognitive profiles seen in AD are thought to reflect the relative distribution of neuropathology, and have been associated with distinct functional neuroimaging patterns⁸⁵⁻⁹⁰.

Wide metacognitive variability across individuals in the mildest stages of AD may signal dissociations in the underlying distribution of disease, although the relevant neural networks are not yet known. Most neuropsychological studies in AD have suggested a relationship between frontal lobe functioning and awareness^{21, 22}, and this relationship has been supported by several functional neuroimaging studies pointing to decreased activation in areas of the right frontal lobe^{6, 91, 92}. This is largely consistent with research in stroke^{67, 71, 73, 93-96}, dementia^{6, 97-100}, and MCI¹⁰¹ that has implicated the importance of right rather than left hemisphere areas in supporting self-awareness related to a wide range of cognitive and behavioral changes. For example, unawareness of hemiplegia is a form of anosognosia frequently reported in the context of right rather than left parietal lesions^{71, 93, 102}. Conversely, altered awareness of one's behavior and social appropriateness is a key feature of the behavioral disorder subtype of frontotemporal dementia (FTD) that may reflect a critical involvement of the right prefrontal cortex in contrast to the primary aphasia syndrome associated with predominantly left hemisphere pathology¹⁰³. Other syndromes of disordered self-awareness such as reduplication and misidentification have been tied to right-sided frontal and parietal injuries^{69, 104-106}, and a recent case study of a commissurotized patient found a critical role for the right hemisphere in self-voice recognition¹⁰⁷. Interestingly, the right hemisphere's role in evaluating aspects of the self has been echoed in studies of healthy adults as well (e.g., face recognition)¹⁰⁸⁻¹¹¹.

Prigatano has insightfully suggested that awareness for specific abilities may be supported by the cortical areas implicated in production of the relevant skill such that the parietal lobes play a primary role in mediating self-awareness for spatial abilities while the prefrontal cortex may be critical to aspects of social self-awareness^{73, 112}. This notion of domain relevant anosognosia might predict a role for the medial temporal

cortex, right greater than left, in supporting awareness for memory deficits. Interestingly, results from the only neuropathological study evaluating anosognosia in AD are fairly consistent with this idea, pointing to decreased cell count in the prosubiculum region of the right hippocampus in unaware patients¹⁰⁰. Current results also lend preliminary support to the idea that the decreased awareness may be related to right rather than left medial temporal integrity. Specifically, while the UAD and AAD groups achieved comparable scores on several measures of cognition and memory, the UAD group scored well below the AAD group on a measure of nonverbal memory. The nonverbal episodic memory task required participants to pair five designs with specific locations in a 4 X 2 grid, exerting demands on the formation of new associations, a function believed to be specifically dependent on hippocampal functioning. Unfortunately, the current study did not include a comprehensive neuropsychological battery; however, ongoing data collection is dedicated to evaluating a broad range of neuropsychological abilities related to both right and left hemisphere functioning. Future neuroimaging studies are needed to more fully investigate the neuroanatomic profile associated with impaired awareness for memory loss, and objective evaluation of memory monitoring may facilitate such investigations.

This study has several limitations. Notably, gamma cannot be calculated when there is no variability in ratings or accuracy, and increasing the number of items on the test will likely introduce the needed variability. This was particularly an issue for the nonverbal episodic task. Further, gamma is sensitive to individual items, and increasing the number of test items may stabilize this variable; this modification has been introduced into a follow up study. Nonetheless, verbal episodic gamma scores correlated significantly with clinical ratings of awareness made prior to all cognitive and metacognitive testing, but with no other clinical or demographic variables. Further, calibration scores and implementation of the Memory for Past Test heuristic differed as a function of awareness.

The current study indicates that with further refinement, the verbal episodic memory monitoring task has the potential to serve as a quantitative measure of metamemory in AD, and may offer important insight into the manner in which metamemory breaks down. Additional work is being conducted to determine the stability of performance on these measures over time, as well as to refine the metacognitive tasks. The utility of a metamemory task as a research instrument is specifically driven by objective administration and scoring. Such psychometric qualities increase the likelihood of

reliable measurement across time in longitudinal analyses as well as across studies. The current study suggests that the current task is a promising approach to assessing metamemory in AD and relating such abilities to disease variables. Further, for practical purposes, our results support the use of ordinal clinical ratings as efficient and qualitative means of characterizing patients' level of awareness of memory loss in the clinic. Finally, metacognitive tasks such as the one included in this paper may have important implications for the diagnosis of Mild Cognitive Impairment (MCI) and we are currently collecting data to examine this issue. Individuals with MCI are considered to be more aware of memory difficulties than individuals with AD; however, this is directly related to the diagnostic criteria for this condition which require subjective memory complaint by either the individual or an informant¹¹³. Studies have begun to document that the heterogeneity in awareness in early AD is mirrored in individuals who meet the memory criteria for MCI^{101, 114}. Longitudinal studies are required to fully appreciate the spectrum of awareness for memory loss that exists at the transition from healthy aging to MCI and to AD.

Table 1

*Demographic and Cognitive Scores in Healthy Elders compared to Entire AD Group
Mean (SD)*

	HE	AD	F	df	p
Age	73.70 (6.85)	76.96 (7.12)	2.35	1, 42	.13
Education	15.70 (2.47)	15.50 (3.51)	0.05	1, 42	.83
% Female	50	50	--	1	1.00
% Caucasian	95	71	--	1	.05
MMSE (0-30)	28.75 (0.85)	25.13 (2.38)	41.70	1, 42	.00
General Knowledge Total (0-13)	9.20 (2.04)	7.21 (2.52)	8.07	1, 42	.01
Verbal Episodic Total (0-20)	18.00 (4.17)	10.39 (5.44)	25.87	1, 37	.00
Verbal Episodic Delay (0-5)	4.60 (1.09)	2.48 (1.40)	29.05	1, 39	.00
Nonverbal Episodic Total (0-20)	18.90 (2.29)	9.64 (5.98)	42.26	1, 40	.00

Note. MMSE = Mini-Mental State Examination (Folstein, Folstein, & McHugh, 1975);

Table 2
Demographic and Cognitive Scores in Aware versus Unaware AD Groups
Mean (SD)

	AAD	UAD	F	df	p
Age	76.87 (6.41)	77.11 (8.58)	0.01	1, 22	.94
Education	14.67 (3.56)	16.89 (3.14)	2.39	1, 22	.14
% Female	60	33	----	1	.40
% Caucasian	67	78	----	1	.67
MMSE (0-30)	25.73 (2.09)	24.11 (2.62)	2.81	1, 22	.11
General Knowledge Total (0-13)	7.27 (2.46)	7.11 (2.76)	0.02	1, 22	.89
Verbal Episodic Total (0-20)	11.27 (5.50)	8.75 (5.28)	1.12	1, 21	.30
Verbal Episodic Delay (0-5)	2.50 (1.45)	2.43 (1.40)	0.01	1, 19	.92
Nonverbal Episodic Total (0-20)	11.67 (5.85)	5.29 (3.59)	6.98	1, 20	.02

Note. AAD = Aware AD group; UAD = Unaware AD group; MMSE = Mini-Mental State Examination (Folstein, Folstein, & McHugh, 1975).

Table 3
Correlational Analyses Relating Clinical Ratings of Awareness to Metacognitive, Cognitive, and Demographic variables in Alzheimer's Disease

	CRA		Verbal Episodic Gamma		NV Global Calibration	
	<i>r</i>	<i>p</i>	<i>r</i>	<i>p</i>	<i>r</i>	<i>p</i>
<u>Metacognitive</u>						
Verbal Episodic Gamma	.46	.03	NA	NA	.00	.99
Verbal Episodic Global Calibration	-.30	.19	.11	.63	.73	.00
Verbal Episodic Item Calibration	-.18	.44	.00	.99	.00	.99
Verbal Episodic Delayed Calibration	-.16	.49	-.14	.56	.53	.02
NV Episodic Gamma	-.03	.90	-.22	.38	.22	.40
NV Episodic Global Calibration	-.50	.03	.00	.99	NA	NA
NV Episodic Item Calibration	-.41	.08	-.13	.60	.63	.01
General Knowledge Gamma	.30	.15	.36	.09	-.28	.23
General Knowledge Calibration	-.12	.59	-.27	.23	.25	.30
<u>Cognitive</u>						
MMSE	.29	.18	-.02	.94	-.30	.21
Verbal Episodic Total	.06	.79	-.05	.83	-.24	.32
Verbal Episodic Delay	-.04	.85	.04	.86	-.24	.30
Nonverbal Episodic Total	.31	.16	.08	.72	-.78	.00
General Knowledge Total	.04	.86	.13	.55	.25	.29
<u>Demographic</u>						
Age	-.01	.96	-.13	.55	-.16	.49
Education	-.19	.37	-.21	.34	.40	.08
<u>Correlations controlling for MMSE</u>						
Verbal Episodic Gamma	.48	.02				
NV Episodic Global Calibration	-.53	.02				

Table 4
Mean Metacognitive Scores by Group
Mean (SD)

	HE	AAD	UAD
Verbal Episodic Gamma	0.69 (0.66)	0.76 (0.36)	0.28 (0.83) [^]
Verbal Episodic Global Calibration	-0.03 (0.07)	-0.06 (0.28)	0.15 (0.23) ^{*,^}
Verbal Episodic Item Calibration	0.00 (0.07)	0.06 (.16)	0.12 (0.14) [*]
Verbal Episodic Delayed Calibration	-0.11 (0.18)	-0.10 (0.28)	0.04 (0.27)
NV Episodic Gamma	0.55 (0.78)	0.39 (0.73)	0.63 (0.47)
NV Episodic Global Calibration	-0.23 (0.36)	-0.07 (0.33)	0.37 (0.16) ^{*,^}
NV Episodic Item Calibration	0.03 (0.12)	0.10 (0.14)	0.30 (0.30) ^{*,^}
General Knowledge Gamma	0.57 (0.48)	0.62 (0.27)	0.43 (0.40)
General Knowledge Calibration	-0.02 (0.13)	-0.05 (0.15)	0.05 (0.11)

Note. * Indicates performance is significantly different from healthy elders ($p < .05$).

[^] Indicates performance is significantly different within AD groups.


Table 5

Use of Memory for Past Test Heuristic

	Verbal Episodic			Nonverbal Episodic		
	<i>M</i> (<i>SD</i>)	<i>t</i>	<i>p</i>	<i>M</i> (<i>SD</i>)	<i>t</i>	<i>p</i>
Healthy Elders						
Trial 2	0.40 (0.68)	2.63	.02	0.00 (1.21)	0.00	1.0
Trial 3	0.33 (0.78)	1.48	.17	0.00 (1.63)	0.00	1.0
Trial 4	0.00 (0.00)	NA	NA	0.33 (0.82)	1.00	.36
Aware AD						
Trial 2	0.43 (1.27)	1.32	.21	-0.46 (1.14)	-1.45	.17
Trial 3	-0.40 (1.76)	-0.88	.40	0.00 (1.49)	0.00	1.00
Trial 4	-0.21 (1.29)	-0.56	.59	-0.08 (1.72)	-0.17	.87
Unaware AD						
Trial 2	0.83 (1.27)	1.96	.09	1.25 (1.28)	2.76	.03
Trial 3	0.81 (1.25)	1.83	.11	1.86 (1.07)	4.60	.00
Trial 4	1.00 (0.75)	2.37	.05	2.17 (0.75)	7.05	.00

Note. Results are from one-sample t-tests conducted in each group to determine if MPT values were significantly different from zero. Positive mean values represent predictions that were higher than the accuracy achieved on the former trial and negative values represent predictions that were lower than the accuracy achieved on the former trial.

Figure 1: Item-Level Calibration Scores for Trials Blocks on the Verbal Episodic Task



REFERENCES

1. Hyman BT, Van Hoesen GW, Damasio AR, Barnes CL. Alzheimer's disease: cell-specific pathology isolates the hippocampal formation. *Science*. Sep 14 1984;225(4667):1168-1170.
2. Braak H, Braak E. Neuropathological staging of Alzheimer-related changes. *Acta Neuropathol (Berl)*. 1991;82(4):239-259.
3. Squire LR. Memory and the hippocampus: a synthesis from findings with rats, monkeys, and humans. *Psychol Rev*. Apr 1992;99(2):195-231.
4. Storandt M, Botwinick J, Danziger WL, Berg L, Hughes CP. Psychometric differentiation of mild senile dementia of the Alzheimer type. *Arch Neurol*. May 1984;41(5):497-499.
5. Neary D, Snowden JS, Bowen DM, Sims NR, Mann DM, Benton JS, Northen B, Yates PO, Davison AN. Neuropsychological syndromes in presenile dementia due to cerebral atrophy. *J Neurol Neurosurg Psychiatry*. Feb 1986;49(2):163-174.
6. Reed BR, Jagust WJ, Coulter L. Anosognosia in Alzheimer's disease: relationships to depression, cognitive function, and cerebral perfusion. *J Clin Exp Neuropsychol*. Mar 1993;15(2):231-244.
7. Smith CA, Henderson VW, McCleary CA, Murdock GA, Buckwalter JG. Anosognosia and Alzheimer's disease: the role of depressive symptoms in mediating impaired insight. *J Clin Exp Neuropsychol*. Aug 2000;22(4):437-444.
8. Hunt L, Morris JC, Edwards D, Wilson BS. Driving performance in persons with mild senile dementia of the Alzheimer type. *J Am Geriatr Soc*. Jul 1993;41(7):747-752.
9. Cotrell V, Wild K. Longitudinal study of self-imposed driving restrictions and deficit awareness in patients with Alzheimer disease. *Alzheimer Dis Assoc Disord*. Jul-Sep 1999;13(3):151-156.
10. Koltai DC, Welsh-Bohmer KA, Schmechel DE. Influence of anosognosia on treatment outcome among dementia patients. *Neuropsychological Rehabilitation*. 2001;11:455-475.
11. Clare L, Wilson BA, Carter G, Roth I, Hodges JR. Awareness in early-stage Alzheimer's disease: relationship to outcome of cognitive rehabilitation. *J Clin Exp Neuropsychol*. Apr 2004;26(2):215-226.
12. Seltzer B, Vasterling JJ, Yoder JA, Thompson KA. Awareness of deficit in Alzheimer's disease: relation to caregiver burden. *Gerontologist*. Feb 1997;37(1):20-24.
13. Spitznagel M, Gunstad J, Carvahlo JO, Brown LB, Davis JD, Tremont G. Impaired Awareness in Dementia Patients Predicts Caregiver Burden [abstract]. *Journal of the International Neuropsychological Society*. 2006;12(S1):51.
14. Agnew SK, Morris RG. The heterogeneity of anosognosia for memory impairment in Alzheimer's disease: A review of the literature and a proposed model. *Aging and Mental Health*. 1998;2:9-15.

15. Morris R, Hannesdottir K. Loss of Awareness in Alzheimer's Disease. In: Morris R, Becker J, eds. *Cognitive Neuropsychology of Alzheimer's Disease*. Oxford, UK: Oxford University Press; 2004:275-296.
16. Cosentino S, Stern Y. Metacognitive theory and assessment in dementia: do we recognize our areas of weakness? *J Int Neuropsychol Soc*. Nov 2005;11(7):910-919.
17. Clare L. Awareness in early-stage Alzheimer's disease: a review of methods and evidence. *Br J Clin Psychol*. Jun 2004;43(Pt 2):177-196.
18. Zanetti O, Vallotti, B., Frisoni, G.B., Geroldi, C., Bianchetti, A., Pasqualetti, P., Trabucchi, M. Insight in dementia: when does it occur? Evidence for a nonlinear relationship between insight and cognitive status. *Journal of Gerontology: Psychological Sciences*. 1999;54(2):100-106.
19. McDaniel KD, Edland, S.D., Heyman, A. Relationship between level of insight and severity of dementia in Alzheimer disease. CERAD Clinical Investigators. Consortium to Establish a Registry for Alzheimer's Disease. *Alzheimer Dis Assoc Disord*. 1995;9(2):101-104.
20. Derouesne C, Thibault, S., Lagha-Pierucci, S., Baudouin-Madec, V., Ancrì, D., Lacomblez, L. Decreased awareness of cognitive deficits inpatients with mild dementia of the Alzheimer type. *International Journal of Geriatric Psychiatry*. 1999;14:1019-1030.
21. Mangone CA, Hier DB, Gorelick PB, Ganellen RJ, Langenberg P, Boarman R, Dollear WC. Impaired insight in Alzheimer's disease. *J Geriatr Psychiatry Neurol*. Oct-Dec 1991;4(4):189-193.
22. Michon A, Deweer B, Pillon B, Agid Y, Dubois B. Relation of anosognosia to frontal lobe dysfunction in Alzheimer's disease. *J Neurol Neurosurg Psychiatry*. Jul 1994;57(7):805-809.
23. DeBettignies BH, Mahurin RK, Pirozzolo FJ. Insight for impairment in independent living skills in Alzheimer's disease and multi-infarct dementia. *J Clin Exp Neuropsychol*. Mar 1990;12(2):355-363.
24. Kotler-Cope S, Camp CJ. Anosognosia in Alzheimer disease. *Alzheimer Dis Assoc Disord*. Spring 1995;9(1):52-56.
25. Saravanan B, Jacob KS, Prince M, Bhugra D, David AS. Culture and insight revisited. *Br J Psychiatry*. Feb 2004;184:107-109.
26. White R, Bebbington P, Pearson J, Johnson S, Ellis D. The social context of insight in schizophrenia. *Soc Psychiatry Psychiatr Epidemiol*. Nov 2000;35(11):500-507.
27. Sussman LK, ed. *The role of culture in definitions, interpretations, and management of illness*. Mahwah, NJ: Lawrence Erlbaum Associates; 2004. Gielen UP, Fish JM, Draguns JG, eds. *Handbook of culture, therapy, and healing*.
28. Prigatano GP, Weinstein EA. Edwin A. Weinstein's contributions to neuropsychological rehabilitation. *Neuropsychological Rehabilitation*. 1996;6:305-326.
29. Ownsworth T, Clare L, Morris R. An integrated biopsychosocial approach to understanding awareness deficits in Alzheimer's disease and brain injury. *Neuropsychological Rehabilitation*. 2006;16(4):415-438.

30. Clare L. The construction of awareness in early-stage Alzheimer's disease: a review of concepts and models. *Br J Clin Psychol*. Jun 2004;43(Pt 2):155-175.
31. Markova IS, Clare L, Wang M, Romero B, Kenny G. Awareness in dementia: conceptual issues. *Aging Ment Health*. Sep 2005;9(5):386-393.
32. Metcalfe J, Shimamura, A.P., ed. *Metacognition: Knowing about Knowing*. London: The MIT Press; 1994.
33. Hart JT. Memory and the feeling of knowing experience. *Journal of Educational Psychology*. 1965;56:208-216.
34. Nelson TO, Narens, L. Norms of 300 general information questions: Accuracy of recall and feeling-of-knowing ratings. *Journal of verbal learning and verbal behavior*. 1980;19:338-368.
35. Leonesio RJ, Nelson TO. Do different metamemory judgments tap the same underlying aspects of memory? *J Exp Psychol Learn Mem Cogn*. May 1990;16(3):464-467.
36. Schacter DL. Feeling-of-knowing in episodic memory. *Journal of Experimental Psychology: Learning, Memory, and Cognition*. 1983;9:39-54.
37. Nelson TO, Narens L. Why investigate metacognition? In: Metcalfe J, Shimamura A, eds. *Metacognition: Knowing about Knowing*. Cambridge, MA: The MIT Press; 1994.
38. Koriat A, Sheffer L, Ma'ayan H. Comparing objective and subjective learning curves: judgments of learning exhibit increased underconfidence with practice. *J Exp Psychol Gen*. Jun 2002;131(2):147-162.
39. Finn B, Metcalfe J. The role of memory for past test in the underconfidence with practice effect. *Journal of Experimental Psychology: Learning, Memory, and Cognition*. 2007;33(1):238-244.
40. Butterfield EC, Nelson TO, Peck V. Developmental aspects of the feeling of knowing. *Developmental Psychology*. 1988;24:654-663.
41. Connor LT, Dunlosky J, Hertzog C. Age-related differences in absolute but not relative metamemory accuracy. *Psychol Aging*. Mar 1997;12(1):50-71.
42. Pappas BA, Sunderland T, Weingartner HM, Vitello B, Martinson H, Putnam K. Alzheimer's disease and feeling-of-knowing for knowledge and episodic memory. *Journal of Gerontology: Psychological Sciences*. 1992;47(3):159-164.
43. Hertzog C, Dunlosky J, Powell-Moman A, Kidder DP. Aging and monitoring associative learning: is monitoring accuracy spared or impaired? *Psychol Aging*. Jun 2002;17(2):209-225.
44. Dunlosky J, Hertzog C. Updating knowledge about encoding strategies: a componential analysis of learning about strategy effectiveness from task experience. *Psychol Aging*. Sep 2000;15(3):462-474.
45. Lovelace EA, Marsh GR. Prediction and evaluation of memory performance by young and old adults. *J Gerontol*. Mar 1985;40(2):192-197.
46. Shaw RJ, Craik FI. Age differences in predictions and performance on a cued recall task. *Psychol Aging*. Jun 1989;4(2):131-135.

47. Rabinowitz JC, Ackerman BP, Craik FI, Hinchley JL. Aging and metamemory: the roles of relatedness and imagery. *J Gerontol*. Nov 1982;37(6):688-695.
48. Souchay C, Isingrini M, Clarys D, Taconnat L, Eustache F. Executive functioning and judgment-of-learning versus feeling-of-knowing in older adults. *Exp Aging Res*. Jan-Mar 2004;30(1):47-62.
49. Souchay C, Isingrini M, Espagnet L. Aging, episodic memory feeling-of-knowing, and frontal functioning. *Neuropsychology*. Apr 2000;14(2):299-309.
50. Vilkki J, Servo A, Surma-aho O. Word list learning and prediction of recall after frontal lobe lesions. *Neuropsychology*. Apr 1998;12(2):268-277.
51. Janowsky JS, Shimamura, A.P., Squire, L.R. Memory and metamemory: Comparisons between patients with frontal lobe lesions and amnesic patients. *Psychobiology*. 1989;17:3-11.
52. Shimamura AP, Squire LR. Memory and metamemory: a study of the feeling-of-knowing phenomenon in amnesic patients. *J Exp Psychol Learn Mem Cogn*. Jul 1986;12(3):452-460.
53. Backman L, Lipinska B. Monitoring of general knowledge: evidence for preservation in early Alzheimer's disease. *Neuropsychologia*. Apr 1993;31(4):335-345.
54. Lipinska B, Backman L. Feeling-of-knowing in fact retrieval: further evidence for preservation in early Alzheimer's disease. *J Int Neuropsychol Soc*. Jul 1996;2(4):350-358.
55. Moulin CJ, Perfect TJ, Jones RW. Evidence for intact memory monitoring in Alzheimer's disease: metamemory sensitivity at encoding. *Neuropsychologia*. 2000;38(9):1242-1250.
56. McGlynn SM, Kaszniak, A.W. When metacognition fails: Impaired awareness of deficit in Alzheimer's disease. *Journal of Cognitive Neuroscience*. 1991;3:183-189.
57. Moulin CJ, Perfect TJ, Jones RW. The effects of repetition on allocation of study time and judgements of learning in Alzheimer's disease. *Neuropsychologia*. 2000;38(6):748-756.
58. Ansell EL, Bucks RS. Mnemonic anosognosia in Alzheimer's disease: A test of Agnew and Morris (1998). *Neuropsychologia*. 2006;44:1095-1102.
59. Lopez OL, Becker JT, Somsak D, Dew MA, DeKosky ST. Awareness of cognitive deficits and anosognosia in probable Alzheimer's disease. *Eur Neurol*. 1994;34(5):277-282.
60. Souchay C, Isingrini M, Gil R. Alzheimer's disease and feeling-of-knowing in episodic memory. *Neuropsychologia*. 2002;40(13):2386-2396.
61. Moulin CJ. Sense and Sensitivity: Metacognition in Alzheimer's Disease. In: Perfect TJ, Schwartz BL, eds. *Applied Metacognition*. Cambridge, UK: Cambridge University Press; 2002.
62. Reed BR, Seab JP, Jagust WJ. Dementia severity, memory impairment, and awareness of memory loss in Alzheimer's disease. *Journal of Clinical and Experimental Neuropsychology*. 1992;14:21.

63. Deckel AW, Morrison, D. Evidence of a neurologically based "denial of illness" in patients with Huntington's disease. *Archives of Clinical Neuropsychology*. 1996;11(4):295-302.
64. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res*. Dec 1975;12(3):189-198.
65. Adams W, Sheslow D. *WRAML Manual*. Wilmington, DE: Jastak Associates; 1990.
66. Nelson TO, Narens, L. A comparison of current measures of the accuracy of feeling-of-knowing predictions. *psychological Bulletin*. 1984;95:109-133.
67. Bisiach E, Vallar G, Perani D, Papagno C, Berti A. Unawareness of disease following lesions of the right hemisphere: anosognosia for hemiplegia and anosognosia for hemianopia. *Neuropsychologia*. 1986;24(4):471-482.
68. Cutting J. Study of anosognosia. *Journal of Neurology, Neurosurgery and Psychiatry*. 1978;41:548-555.
69. Feinberg TE. *Altered egos: How the brain creates the self*. New York: Oxford University Press; 2001.
70. Geschwind N. Disconnexion syndromes in animals and man. *Brain*. 1965;88:237-294.
71. Heilman KM, Watson, R.T., Valenstein, E. Neglect and related disorders. In: Heilman KM, & Valenstein, E., ed. *Clinical Neuropsychology*. 3rd ed. New York, NY: Oxford University Press; 1993:279-336.
72. Marcel AJ, Tegner R, Nimmo-Smith I. Anosognosia for plegia: specificity, extension, partiality and disunity of bodily unawareness. *Cortex*. Feb 2004;40(1):19-40.
73. Prigatano GP, Schacter DL. *Awareness of Deficit after Brain Injury*. New York: Oxford University Press; 1991.
74. Ramachandran VS. Anosognosia in parietal lobe syndrome. *Conscious Cogn*. Mar 1995;4(1):22-51.
75. Schacter DL. Toward a Cognitive Neuropsychology of Awareness: Implicit Knowledge and Anosognosia. *Journal of Clinical and Experimental Neuropsychology*. 1990;12(1):155-178.
76. Shimamura AP. Toward a cognitive neuroscience of metacognition. *Conscious Cogn*. Jun 2000;9(2 Pt 1):313-323; discussion 324-316.
77. Vuilleumier P. Anosognosia: the neurology of beliefs and uncertainties. *Cortex*. Feb 2004;40(1):9-17.
78. Weinstein E, Kahn R. *Denial of Illness*. Springfield, Illinois: Charles C Thomas; 1955.
79. Kaszniak AW, Zak M. On the neuropsychology of metamemory: Contributions from the study of amnesia and dementia. *Learning and Individual Differences*. 1996;8(4):355-381.
80. Caine D, Hodges JR. Heterogeneity of semantic and visuospatial deficits in early Alzheimer's disease. *Neuropsychology*. Apr 2001;15(2):155-164.

81. Haxby JV, Grady CL, Koss E, Horwitz B, Schapiro M, Friedland RP, Rapoport SI. Heterogeneous anterior-posterior metabolic patterns in dementia of the Alzheimer type. *Neurology*. Dec 1988;38(12):1853-1863.
82. Cummings JL. Cognitive and behavioral heterogeneity in Alzheimer's disease: seeking the neurobiological basis. *Neurobiol Aging*. Nov-Dec 2000;21(6):845-861.
83. Mann UM, Mohr E, Gearing M, Chase TN. Heterogeneity in Alzheimer's disease: progression rate segregated by distinct neuropsychological and cerebral metabolic profiles. *J Neurol Neurosurg Psychiatry*. Oct 1992;55(10):956-959.
84. Strite D, Massman PJ, Cooke N, Doody RS. Neuropsychological asymmetry in Alzheimer's disease: verbal versus visuoconstructional deficits across stages of dementia. *J Int Neuropsychol Soc*. Sep 1997;3(5):420-427.
85. Aharon-Peretz J, Israel O, Goldsher D, Peretz A. Posterior cortical atrophy variants of Alzheimer's disease. *Dement Geriatr Cogn Disord*. Nov-Dec 1999;10(6):483-487.
86. Binetti G, Magni E, Padovani A, Cappa SF, Bianchetti A, Trabucchi M. Executive dysfunction in early Alzheimer's disease. *J Neurol Neurosurg Psychiatry*. Jan 1996;60(1):91-93.
87. Galton CJ, Patterson K, Xuereb JH, Hodges JR. Atypical and typical presentations of Alzheimer's disease: a clinical, neuropsychological, neuroimaging and pathological study of 13 cases. *Brain*. Mar 2000;123 Pt 3:484-498.
88. Johnson JK, Head E, Kim R, Starr A, Cotman CW. Clinical and pathological evidence for a frontal variant of Alzheimer disease. *Arch Neurol*. Oct 1999;56(10):1233-1239.
89. Kanne SM, Balota DA, Storandt M, McKeel DW, Jr., Morris JC. Relating anatomy to function in Alzheimer's disease: neuropsychological profiles predict regional neuropathology 5 years later. *Neurology*. Apr 1998;50(4):979-985.
90. Lambon Ralph MA, Patterson K, Graham N, Dawson K, Hodges JR. Homogeneity and heterogeneity in mild cognitive impairment and Alzheimer's disease: a cross-sectional and longitudinal study of 55 cases. *Brain*. Nov 2003;126(Pt 11):2350-2362.
91. Vogel A, Hasselbalch SG, Gade A, Ziebell M, Waldemar G. Cognitive and functional neuroimaging correlate for anosognosia in mild cognitive impairment and Alzheimer's disease. *Int J Geriatr Psychiatry*. Mar 2005;20(3):238-246.
92. Starkstein SE, Vazquez S, Migliorelli R, Teson A, Sabe L, Leiguarda R. A single-photon emission computed tomographic study of anosognosia in Alzheimer's disease. *Arch Neurol*. Apr 1995;52(4):415-420.
93. Critchley M. *The Parietal Lobes*. New York: Hafner Press; 1953.
94. Heilman KM, Bowers, D., Valenstein, E. *Emotional Disorders Associated with Neurological Diseases*. 3rd ed. New York: Oxford University Press; 1993.

95. Schnyer DM, Verfaellie M, Alexander MP, LaFleche G, Nicholls L, Kaszniak AW. A role for right medial prefrontal cortex in accurate feeling-of-knowing judgements: evidence from patients with lesions to frontal cortex. *Neuropsychologia*. 2004;42(7):957-966.
96. Rainville C, Giroire JM, Periot M, Cuny E, Mazaux JM. The impact of right subcortical lesions on executive functions and spatio-cognitive abilities: a case study. *Neurocase*. Aug 2003;9(4):356-367.
97. Miller BL, Chang L, Mena I, Boone K, Lesser IM. Progressive right frontotemporal degeneration: clinical, neuropsychological and SPECT characteristics. *Dementia*. May-Aug 1993;4(3-4):204-213.
98. Edwards-Lee T, Miller BL, Benson DF, Cummings JL, Russell GL, Boone K, Mena I. The temporal variant of frontotemporal dementia. *Brain*. Jun 1997;120 (Pt 6):1027-1040.
99. Rankin KP, Glenn S, Stanley CM, Allison SA, Kramer JH, Miller BL. Specific right frontal structures mediate social self-monitoring in dementia. *Journal of the International Neuropsychological Society*. 2006;12(S1):121.
100. Marshall GA, Kaufer DI, Lopez OL, Rao GR, Hamilton RL, DeKosky ST. Right prosubiculum amyloid plaque density correlates with anosognosia in Alzheimer's disease. *J Neurol Neurosurg Psychiatry*. Oct 2004;75(10):1396-1400.
101. Adair JC, Acothley R, Knoefel JF. White Matter Abnormalities predict symptom awareness in Mild Cognitive Impairment [abstract]. *Journal of the International Neuropsychological Society*. 2006;12(S1):253.
102. Young GW. *Functions of the right cerebral hemisphere*. London: Academic Press; 1983.
103. Grossman M. Frontotemporal dementia: A review. *Journal of the International Neuropsychological Society*. 2000;8:566-583.
104. Feinberg TE, Eaton LA, Roane DM, Giacino JT. Multiple fregoli delusions after traumatic brain injury. *Cortex*. Jun 1999;35(3):373-387.
105. Feinberg TE, Roane DM, Ali J. Illusory limb movements in anosognosia for hemiplegia. *J Neurol Neurosurg Psychiatry*. Apr 2000;68(4):511-513.
106. Feinberg TE, Roane DM, Kwan PC, Schindler RJ, Haber LD. Anosognosia and visuoverbal confabulation. *Arch Neurol*. May 1994;51(5):468-473.
107. Rosa C, Belin P, Keenan JP, Lassonde M. Self-Voice Recognition: Greater Implication of the Right Hemisphere in a Commissurotomized Patient [abstract]. *Journal of the International Neuropsychological Society*. 2006;12(S1):250.
108. Fink GR, Markowitsch HJ, Reinkemeier M, Bruckbauer T, Kessler J, Heiss WD. Cerebral representation of one's own past: neural networks involved in autobiographical memory. *J Neurosci*. Jul 1 1996;16(13):4275-4282.
109. Keenan JP, Freund S, Hamilton RH, Ganis G, Pascual-Leone A. Hand response differences in a self-face identification task. *Neuropsychologia*. 2000;38(7):1047-1053.
110. Keenan JP, McCutcheon B, Freund S, Gallup GG, Jr., Sanders G, Pascual-Leone A. Left hand advantage in a self-face recognition task. *Neuropsychologia*. Nov 1999;37(12):1421-1425.

111. Platek SM, Keenan JP, Gallup GG, Jr., Mohamed FB. Where am I? The neurological correlates of self and other. *Brain Res Cogn Brain Res*. Apr 2004;19(2):114-122.
112. Prigatano GP. Disturbances in self-awareness after traumatic brain injury. In: Prigatano GP, Schacter DL, eds. *Awareness of Deficit After Brain Injury*. New York: Oxford University Press; 1991:111-126.
113. Petersen RC. Mild cognitive impairment as a diagnostic entity. *J Intern Med*. Sep 2004;256(3):183-194.
114. Vogel A, Stokholm J, Gade A, Andersen BB, Hejl AM, Waldemar G. Awareness of deficits in mild cognitive impairment and Alzheimer's disease: do MCI patients have impaired insight? *Dement Geriatr Cogn Disord*. 2004;17(3):181-187.